

THE FUNCTOR $A^{\min}$ FOR $(p-1)$ -CELL COMPLEXES AND EHP SEQUENCES

J. WU[†]

ABSTRACT. Let X be a co- H -space of $(p-1)$ -cell complex with all cells in even dimensions. Then the loop space ΩX admits a retract $\bar{A}^{\min}(X)$ that is the evaluation of the functor $\bar{A}^{\min}$ on X . In this paper, we determine the homology $H_*(\bar{A}^{\min}(X))$ and give the EHP sequence for the spaces $\bar{A}^{\min}(X)$.

1. INTRODUCTION

The algebraic functor $A^{\min}$ was introduced in [16] arising from the question on the naturality of the classical Poincaré-Birkhoff-Witt isomorphism. For any *ungraded* module V , $A^{\min}(V)$ is defined to be the *smallest functorial coalgebra retract* of $T(V)$ containing V . Then the functor $A^{\min}$ extends canonically to the cases when V is any graded module. (See [16] for details.) The functor $A^{\min}$ admits the tensor-length decomposition with

$$A^{\min}(V) = \bigoplus_{n=0}^{\infty} A_n^{\min}(V),$$

where $A_n^{\min}(V) = A^{\min}(V) \cap T_n(V)$ is the homogenous component of $A^{\min}(V)$. By the Functorial Poincaré-Birkhoff-Witt Theorem [16, Theorem 6.5], there exists a functor $B^{\max}$ from (graded) modules to Hopf algebras with the functorial coalgebra decomposition

$$T(V) \cong A^{\min}(V) \otimes B^{\max}(V)$$

for any graded module V . The determination of $A^{\min}(V)$ for general V is equivalent to an open problem in the modular representation theory of the symmetric groups according to [16, Theorem 7.4], which seems beyond the reach of current techniques. Some properties of the algebraic functor $A^{\min}$ have been studied in [7, 23] with applications in homotopy theory [8, 22]. In this article we determine $A^{\min}(V)$ in the special cases when $V_{\text{even}} = 0$ and $\dim V = p-1$.

Denote by $L(V)$ the free Lie algebra generated by V . Write $L_n(V)$ for the n th homogeneous component of $L(V)$. Observe that $[L_s(V), L_t(V)]$ is a submodule of $L_{s+t}(V)$ under the Lie bracket of $L(V)$. Let

$$\bar{L}_n(V) = L_n(V) / \sum_{i=2}^{n-2} [L_i(V), L_{n-i}(V)].$$

Define $\bar{L}_n^k(V)$ recursively by $\bar{L}_n^1(V) = \bar{L}_n(V)$ and $\bar{L}_n^{k+1}(V) = \bar{L}_n(\bar{L}_n^k(V))$.

2000 *Mathematics Subject Classification.* Primary 55P35; Secondary 55Q25, 55P65, 16W30.

Key words and phrases. EHP sequences, functor $A^{\min}$, co- H -spaces, Hopf invariants.

[†] Research is supported in part by the Academic Research Fund of the National University of Singapore R-146-000-101-112.

Theorem 1.1. *Let the ground field $\mathbf{k}$ be of characteristic $p > 2$. Let V be a graded module such that $V_{\text{even}} = 0$ and $\dim V = p - 1$. Then there is an isomorphism of coalgebras*

$$A^{\min}(V) \cong \bigotimes_{k=0}^{\infty} E(\bar{L}_p^k(V)),$$

where $E(W)$ is the exterior algebra generated by W .

An important observation is that

$$\bar{L}_p^k(V)_{\text{even}} = 0 \text{ and } \dim \bar{L}_p^k(V) = p - 1$$

for each $k \geq 0$ provided that $V_{\text{even}} = 0$ and $\dim V = p - 1$. Replace V by $\bar{L}_p(V)$ in the above theorem. Then $A^{\min}(\bar{L}_p(V)) \cong \bigotimes_{k=1}^{\infty} E(\bar{L}_p^k(V))$ with a coalgebra decomposition

$$A^{\min}(V) \cong E(V) \otimes A^{\min}(\bar{L}_p(V)),$$

which indicates the existence of the EHP fibrations by taking the *geometric realization* of the functor $A^{\min}$ and the *Hopf invariants* on the functor $A^{\min}$.

The geometric realization of the (algebraic) functor $A^{\min}$ was studied in [9, 14, 16, 17, 19] for giving the decompositions of the loop spaces of double suspensions of torsion spaces. The most general result so far is given as follows:

Theorem 1.2. [15, Theorem 1.2] *There exist homotopy functors $\bar{A}^{\min}$ and $\bar{B}^{\max}$ from simply connected co- H spaces of finite type to spaces such that for any p -local simply connected co- H space Y the following hold:*

- 1) $\bar{A}^{\min}(Y)$ is a functorial retract of ΩY and so there is a functorial decomposition

$$\Omega Y \simeq \bar{A}^{\min}(Y) \times \bar{B}^{\max}(Y).$$

- 2) On mod p homology the decomposition

$$H_*(\Omega Y) \cong H_*(\bar{A}^{\min}(Y)) \otimes H_*(\bar{B}^{\max}(Y))$$

is with respect to the augmentation ideal filtration.

- 3) On mod p homology the associated bigraded $E^0 H_*(\bar{A}^{\min}(Y))$ is given by

$$E^0 H_*(\bar{A}^{\min}(Y)) = A^{\min}(\Sigma^{-1} \bar{H}_*(Y)).$$

Let X be a path-connected finite complex. Define $b_X = \sum_{q=1}^{\infty} q \dim \bar{H}_q(X; \mathbb{Z}/p)$.

Roughly speaking, b_X is the summation of the dimensions of the cells in X . A direct consequence of Theorems 1.1 and 1.2 is the following homological information:

Corollary 1.3. *Let $p > 2$ and let Y be any p -local simply connected co- H space. Suppose that $\bar{H}_{\text{odd}}(Y) = 0$ and $\dim \bar{H}_*(Y) = p - 1$. Then there is an isomorphism of coalgebras*

$$H_*(\bar{A}^{\min}(Y)) \cong E(\Sigma^{-1} \bar{H}_*(Y)) \otimes \bigotimes_{k=1}^{\infty} E(\Sigma^{\frac{p^k-1}{p-1} b_Y - p^k} \bar{H}_*(Y)),$$

where $\Sigma^{\frac{p^k-1}{p-1} b_Y - p^k} \bar{H}_*(Y) = \bar{L}_p^k(\Sigma^{-1} \bar{H}_*(Y))$. □

Recall that the classical Hopf invariants can be obtained from the suspension splitting of the loop suspensions. For the geometric functor $\bar{A}^{\min}$, the following suspension splitting theorem is a special case of Theorem 4.4.

Theorem 1.4. *Let Y be any p -local simply connected co- H space. Then there is a suspension splitting*

$$\Sigma \bar{A}^{\min}(Y) \simeq \bigvee_{n=1}^{\infty} \bar{A}_n^{\min}(Y)$$

such that

$$\Sigma^{-1} \bar{H}_*(\bar{A}_n^{\min}(Y)) \cong A_n^{\min}(\Sigma^{-1} \bar{H}_*(Y))$$

for each $n \geq 1$. $\square$

Note that each $\bar{A}_n^{\min}(Y)$ is a co- H space because it is a retract of $\Sigma \bar{A}^{\min}(Y)$. From the above theorem, one gets the Hopf invariant H_n defined as the composite

$$\bar{A}^{\min}(Y) \hookrightarrow \Omega \Sigma \bar{A}^{\min}(Y) \simeq \Omega \bigvee_{n=1}^{\infty} \bar{A}_n^{\min}(Y) \xrightarrow{\text{proj}} \Omega \bar{A}_n^{\min}(Y) \xrightarrow{r} \bar{A}^{\min}(\bar{A}_n^{\min}(Y)),$$

where $r: \Omega Z \rightarrow \bar{A}^{\min}(Z)$ is the functorial retraction for any simply connected co- H -space Z . In particular, there is a Hopf invariant

$$H_p: \bar{A}^{\min}(Y) \rightarrow \bar{A}^{\min}(\bar{A}_p^{\min}(Y))$$

for any simply connected co- H space Y . By Lemma 5.3,

$$\bar{A}_p^{\min}(Y) \simeq \Sigma^{b_Y - p + 1} Y$$

provided that $\bar{H}_{\text{odd}}(Y) = 0$ and $\dim \bar{H}_*(Y) = p - 1$.

Theorem 1.5 (EHP Fibration). *Let $p > 2$ and let Y be any p -local simply connected co- H space. Suppose that $\bar{H}_{\text{odd}}(Y) = 0$ and $\dim \bar{H}_*(Y) = p - 1$. Then there is a fibre sequence*

$$\Omega \bar{A}^{\min}(\Sigma^{b_Y - p + 1} Y) \xrightarrow{P} \bar{E}(Y) \xrightarrow{E} \bar{A}^{\min}(Y) \xrightarrow{H_p} \bar{A}^{\min}(\Sigma^{b_Y - p + 1} Y).$$

with the following properties:

- 1) On mod p homology $H_*(\bar{E}(Y)) \cong E(\Sigma^{-1} \bar{H}_*(Y))$ as coalgebras.
- 2) $H_*(\bar{A}^{\min}(Y)) \cong H_*(\bar{E}(Y)) \otimes H_*(\bar{A}^{\min}(\Sigma^{b_Y - p + 1} Y))$ as coalgebras.
- 3) If $f: S^n \rightarrow Y$ is a co- H map such that $f_* \neq 0: \bar{H}_*(S^n) \rightarrow \bar{H}_*(Y)$. Then there is a commutative diagram of fibre sequences

$$\begin{array}{ccccccc} \Omega \bar{A}^{\min}(\Sigma^{b_Y - p + 1} Y) & \xrightarrow{P} & \bar{E}(Y) & \xrightarrow{E} & \bar{A}^{\min}(Y) & \xrightarrow{H_p} & \bar{A}^{\min}(\Sigma^{b_Y - p + 1} Y) \\ \parallel & & \uparrow & & \uparrow & & \parallel \\ \Omega \bar{A}^{\min}(\Sigma^{b_Y - p + 1} Y) & \xrightarrow{P_f} & S^{n-1} & \longrightarrow & B_f & \longrightarrow & \bar{A}^{\min}(\Sigma^{b_Y - p + 1} Y). \end{array}$$

In particular, the map $P: \Omega \bar{A}^{\min}(\Sigma^{b_Y - p + 1} Y) \rightarrow \bar{E}(Y)$ factors through the bottom cell of $\bar{E}(Y)$.

The following theorem gives a general criterion when the EHP fibration splits off. A space X is called to have a *retractile generating complex* C if C is a retract of ΣX with a retraction $r: \Sigma X \rightarrow C$ such that the mod p cohomology $H^*(X)$ is generated by

$$M = \text{Im}(\Sigma^{-1} r^*: \Sigma^{-1} \bar{H}^*(C) \rightarrow \bar{H}^*(X))$$

and $M \cong QH^*(X)$, the set of indecomposables. Recall that a space X is called (stably) *atomic* if any self (stable) map of X inducing isomorphism on the bottom homology is a (stable) homotopy equivalence.

Theorem 1.6. *Let $p > 2$ and let Y be any p -local simply connected co- H space such that Y is stably atomic. Suppose that $\bar{H}_{\text{odd}}(Y) = 0$ and $\dim \bar{H}_*(Y) = p - 1$. Then the following statements are equivalent to each other:*

- 1) *The EHP fibration*

$$\bar{E}(Y) \xrightarrow{E} \bar{A}^{\min}(Y) \xrightarrow{H_p} \bar{A}^{\min}(\Sigma^{b_Y-p+1}Y)$$

splits off.

- 2) $\bar{E}(Y)$ *is an H -space.*
 3) *There exists an H -space X having Y as a retractile generating complex.*
 4) *The map $P: \Omega \bar{A}^{\min}(\Sigma^{b_Y-p+1}Y) \rightarrow \bar{E}(Y)$ is null homotopic.*
 5) *The composite*

$$\Sigma^{b_Y-p-1}Y \hookrightarrow \Omega \bar{A}^{\min}(\Sigma^{b_Y-p+1}Y) \xrightarrow{P} \bar{E}(Y)$$

is null homotopic.

- 6) *There exists a map $g: \Sigma^{b_Y-p}Y \rightarrow \bar{A}^{\min}(Y)$ such that*

$$g_*: H_*(\Sigma^{b_Y-p}Y) \longrightarrow H_*(\bar{A}^{\min}(Y))$$

is a monomorphism on the bottom cells of $\Sigma^{b_Y-p}Y$.

It is a classical question whether the total space of a spherical fibration over a sphere is an H -space localized at an odd prime p . For having a possible H -space structure, the base space and the fibre must be odd dimensional spheres. For $p > 3$, it is well-known that the answer is positive [3, 4]. For $p = 3$, there are examples that do not admit H -space structure. The above theorem gives some properties for studying this question.

The article is organized as follows. In section 2, we study the representation theory on natural coalgebra decompositions of tensor algebras. The proof of Theorem 1.1 is given in Section 3. The geometry of natural coalgebra decompositions of tensor algebras is investigated in Section 4, where Theorem 1.4 is Theorem 4.4. In Section 5, we give the proof of Theorem 1.5. The proof of Theorem 1.6 is given in Section 6.

2. THE FUNCTOR $A^{\min}$ AND THE SYMMETRIC GROUP MODULE $\text{Lie}(n)$

2.1. The Functor $A^{\min}$ on Ungraded Modules. In this subsection, the ground ring is a field $\mathbf{k}$. A coalgebra means a pointed coassociative cocommutative coalgebra. For any module V , the tensor algebra $T(V)$ is a Hopf algebra by requiring V to be primitive. This defines $T: V \mapsto T(V)$ as a functor from *ungraded* modules to *coalgebras*. The functor $A^{\min}: V \mapsto A^{\min}(V)$ is defined to be the smallest coalgebra retract of the functor T with the property that $V \subseteq A^{\min}(V)$. More precisely the functor $A^{\min}$ is defined by the following property:

- (1). $A^{\min}$ is a functor from $\mathbf{k}$ -modules to coalgebras with a natural linear inclusion $V \hookrightarrow A^{\min}(V)$.

- (2). $A^{\min}(V)$ is a natural coalgebra retract of $T(V)$ over V . Namely there exist natural coalgebra transformations $s: A^{\min} \rightarrow T$ and $r: T \rightarrow A^{\min}$ such that the diagram

$$\begin{array}{ccccc} A^{\min}(V) & \xrightarrow{s_V} & T(V) & \xrightarrow{r_V} & A^{\min}(V) \\ \uparrow & & \uparrow & & \uparrow \\ V & \xlongequal{\quad} & V & \xlongequal{\quad} & V \end{array}$$

commutes for any V and $r \circ s = \text{id}_{A^{\min}}$.

- (3). $A^{\min}$ is *minimal* with respect to the above two conditions: if A is any functor from $\mathbf{k}$ -modules to coalgebras with natural linear inclusion $V \hookrightarrow A(V)$ such that $A(V)$ is a natural coalgebra retract of $T(V)$ over V , then $A^{\min}(V)$ is a natural coalgebra retract of $A(V)$ over V .

By the minimal assumption, the functor $A^{\min}$ is unique up to natural equivalence if it exists. The existence of the functor $A^{\min}$ follows from [16, Theorem 4.12].

There is a multiplication on $A^{\min}$ given by the composite

$$A^{\min} \otimes A^{\min} \xrightarrow{s \otimes s} T \otimes T \xrightarrow{\mu} T \xrightarrow{r} A^{\min}$$

and so $A^{\min}$ is a functor from modules to quasi-Hopf algebras, where a quasi-Hopf algebra means a coassociative and cocommutative bi-algebra without assuming the associativity of the multiplication. The multiplication on $A^{\min}$ induces a new natural coalgebra transformation $r'_V: T(V) \rightarrow A^{\min}(V)$ given by

$$r'_V(x_1 \otimes \cdots \otimes x_n) = (((\cdots (x_1 \cdot x_2) \cdot x_3) \cdots) \cdot x_n)$$

for any $x_1 \otimes \cdots \otimes x_n \in V^{\otimes n}$. By the minimal assumption, the composite

$$A^{\min} \xrightarrow{s} T \xrightarrow{r'} A^{\min}$$

is a natural equivalence. Consider $T(V)$ as an $A^{\min}(V)$ -comodule via the map r'_V . According to [16, Proposition 6.1], the cotensor product

$$B^{\max}(V) = \mathbf{k} \square_{A^{\min}(V)} T(V)$$

is natural sub Hopf algebra of $T(V)$ with a natural coalgebra equivalence

$$(2.1) \quad \mathbf{k} \otimes_{B^{\max}(V)} T(V) \cong A^{\min}(V).$$

Together with [16, Lemmas 5.3], there is a natural coalgebra decomposition

$$(2.2) \quad T(V) \cong B^{\max}(V) \otimes A^{\min}(V).$$

By taking tensor length decomposition, we have $B^{\max}(V) = \bigoplus_{n=0}^{\infty} B_n^{\max}(V)$ with $B_n^{\max}(V) = B^{\max}(V) \cap T_n(V)$. Let

$$(2.3) \quad Q^{\max}(V) = QB^{\max}(V)$$

be the indecomposables of $B^{\max}(V)$ with tensor length decomposition $Q^{\max}(V) = \bigoplus_{n=2}^{\infty} Q_n^{\max}(V)$. (Note that $B_1^{\max}(V) = 0$ and so $B^{\max}(V)$ has no nontrivial indecomposable elements of tensor length 1.) According to [16, Section 2], all of the functors $A^{\min}$, $A_n^{\max}$, $B^{\max}$, $Q^{\max}$ and $Q_n^{\max}$ extend canonically for graded modules.

2.2. The Symmetric Group Module $\text{Lie}(n)$. Let the ground ring R be any commutative ring with identity. The module $\text{Lie}^R(n)$, which is simply denoted as $\text{Lie}(n)$ if the ground ring is clear, is defined as follows. Let $\bar{V}$ be a free R -module of rank n with a basis $\{e_i \mid 1 \leq i \leq n\}$. The module γ_n is defined to be the submodule of $\bar{V}^{\otimes n}$ spanned by

$$e_{\sigma(1)} \otimes e_{\sigma(2)} \otimes \cdots \otimes e_{\sigma(n)}$$

for $\sigma \in \Sigma_n$. The module $\text{Lie}(n)$ is defined by

$$\text{Lie}(n) = \gamma_n \cap L_n(\bar{V}) \subseteq \bar{V}^{\otimes n}.$$

The Σ_n -action on γ_n is given by permuting letters $e_1, e_2, \dots, e_n$. Since both γ_n and $L_n(\bar{V})$ are invariant under permutations of the letters, $\text{Lie}(n)$ is an $R(\Sigma_n)$ -submodule of γ_n .

Note that $\text{Lie}(n)$ is the submodule of $L_n(\bar{V})$ spanned by the homogenous Lie elements in which each e_i occurs exactly once. By the Witt formula, $\text{Lie}(n)$ is a free R -module of rank $(n-1)!$. Following from the antisymmetry and the Jacobi identity, $\text{Lie}(n)$ has a basis given by the elements

$$[[e_1, e_{\sigma(2)}], e_{\sigma(3)}], \dots, e_{\sigma(n)}$$

for $\sigma \in \Sigma_{n-1}$. (See [1].) Observe that $\text{Lie}(n)$ is the image of the $R(\Sigma_n)$ -map

$$\beta_n: \gamma_n \rightarrow \gamma_n \quad \beta_n(a_1 \otimes \cdots \otimes a_n) = [[a_1, a_2], \dots, a_n].$$

Thus $\text{Lie}(n)$ can be also regarded as the quotient $R(\Sigma_n)$ -module of γ_n with the projection $\beta_n: \gamma_n \rightarrow \text{Lie}(n)$.

Proposition 2.1. *Let V be any graded projective module and let Σ_n act on $V^{\otimes n}$ by permuting factors in graded sense. Then there is a functorial isomorphism*

$$\text{Lie}(n) \otimes_{R(\Sigma_n)} V^{\otimes n} \cong L_n(V)$$

for any graded module V .

Proof. Clearly the quotient map $\beta_n: V^{\otimes n} = \gamma_n \otimes_{R(\Sigma_n)} V^{\otimes n} \rightarrow L_n(V)$, $x_1 \otimes \cdots \otimes x_n \mapsto [[x_1, x_2], \dots, x_n]$, factors through the quotient $\text{Lie}(n) \otimes_{R(\Sigma_n)} V^{\otimes n}$ and so there is an epimorphism $\text{Lie}(n) \otimes_{R(\Sigma_n)} V^{\otimes n} \twoheadrightarrow L_n(V)$. On the other hand note that $L = \bigoplus_{n=1}^{\infty} \text{Lie}(n) \otimes_{R(\Sigma_n)} V^{\otimes n}$ has the canonical graded Lie algebra structure generated by V . So the inclusion $V \hookrightarrow L$ induces an epimorphism of graded Lie algebras $L(V) \twoheadrightarrow L$. The assertion follows. $\square$

Consider $T_n: V \mapsto V^{\otimes n}$ as a functor from projective (*ungraded*) modules to projective (*ungraded*) modules. Denote by $\text{Hom}(F, F')$ the set of natural linear transformations from a functor F to a functor F' provided that F preserves direct limits. By [7, Lemma 3.8], $\text{Hom}(T_n, T_m) = 0$ if $n \neq m$ and there is an isomorphism of rings

$$(2.4) \quad \theta: \text{End}_{R(\Sigma_n)}(\gamma_n) \longrightarrow \text{End}(T_n, T_n)$$

given by

$$\theta(\phi) = \phi \otimes \text{id}_{V^{\otimes n}}: \gamma_n \otimes_{R(\Sigma_n)} V^{\otimes n} = V^{\otimes n} \longrightarrow \gamma_n \otimes_{R(\Sigma_n)} V^{\otimes n} = V^{\otimes n}$$

for $\phi \in \text{End}_{R(\Sigma_n)}(\gamma_n)$. Replacing γ_n by $\text{Lie}(n)$, we have the morphism of rings $\theta: \text{End}_{R(\Sigma_n)}(\text{Lie}(n)) \longrightarrow \text{End}(L_n)$.

Proposition 2.2. *If $n \neq m$, then $\text{Hom}(L_n, L_m) = 0$. Moreover the map*

$$\theta: \text{End}_{R(\Sigma_n)}(\text{Lie}(n)) \longrightarrow \text{End}(L_n)$$

is an isomorphism.

Proof. Let $\phi: L_n \rightarrow L_m$ be a natural transformation. Then the composite

$$T_n \longrightarrow L_n \xrightarrow{\phi} L_m \hookrightarrow T_m$$

is a natural transformation, which is zero as $\text{Hom}(T_n, T_m) = 0$. Thus $\phi = 0$. For the second statement, clearly θ is a monomorphism. Let $\phi_V: L_n(V) \rightarrow L_n(V)$ be any natural transformation. Let $\bar{V}$ be the free R -module of rank n , which defines γ_n . Consider the commutative diagram

$$\begin{array}{ccccccc} T_n(\bar{V}) & \xrightarrow{\beta_n} & L_n(\bar{V}) & \xrightarrow{\phi_{\bar{V}}} & L_n(\bar{V}) & \hookrightarrow & T_n(\bar{V}) \\ \uparrow & & \uparrow & & \uparrow & & \uparrow \\ \gamma_n & \xrightarrow{\beta_n} & \text{Lie}(n) & \xrightarrow{\phi'} & \text{Lie}(n) & \hookrightarrow & \gamma_n \end{array}$$

where the existence of ϕ' follows from the fact that the composite of the maps in the top row maps γ_n into γ_n and $\text{Lie}(n) = \gamma_n \cap L_n(\bar{V})$. Let V be any ungraded module and let $a_1 \otimes \cdots \otimes a_n$ be any homogenous element in $V^{\otimes n}$. Let $f: \bar{V} \rightarrow V$ be R -linear map such that $f(e_i) = a_i$. By the naturality, there is a commutative diagram

$$\begin{array}{ccccc} \text{Lie}(n) & \hookrightarrow & L_n(\bar{V}) & \xrightarrow{L_n(f)} & L_n(V) \\ \downarrow \phi' & & \downarrow \phi_{\bar{V}} & & \downarrow \phi_V \\ \text{Lie}(n) & \hookrightarrow & L_n(\bar{V}) & \xrightarrow{L_n(f)} & L_n(V) \end{array}$$

Thus

$$\theta(\phi')([a_1, a_2], \dots, a_n) = \phi([a_1, a_2], \dots, a_n)$$

and hence the result. $\square$

Corollary 2.3. *There is a one-to-one correspondence between the decompositions of the functor L_n and the decompositions of $\text{Lie}(n)$ over $R(\Sigma_n)$.* $\square$

A functor from modules to modules Q is called T_n -projective if Q is naturally equivalent to a retract of the functor T_n .

Proposition 2.4. *Let Q be a T_n -projective functor and let $\phi: Q \rightarrow L_n$ be a natural linear transformation. Then there exists a natural linear transformation $\tilde{\phi}: Q \rightarrow T_n$ such that $\phi = \beta_n \circ \tilde{\phi}$.*

Proof. Let $\bar{V}$ be the free R -module of rank n , which defines γ_n . Let $\bar{Q} = \gamma_n \cap Q(\bar{V})$. Since Q is a retract of the functor T_n , $\bar{Q}$ is a summand of γ_n over $R(\Sigma_n)$ and so $\bar{Q}$ is a projective $R(\Sigma_n)$ -module. From the fact that $\text{End}_{R(\Sigma_n)}(\gamma_n) \cong \text{End}(T_n)$,

$$Q(V) = \bar{Q} \otimes_{R(\Sigma_n)} V^{\otimes n}$$

for any module V . By the proof of Proposition 2.2, the map

$$\theta: \text{Hom}_{R(\Sigma_n)}(\bar{Q}, \text{Lie}(n)) \longrightarrow \text{Hom}(Q, L_n), \quad f \mapsto f \otimes \text{id}_{V^{\otimes n}}$$

is an isomorphism. Thus there exists a unique $R(\Sigma_n)$ -map

$$\phi': \bar{Q} \longrightarrow \text{Lie}(n)$$

such that $\theta(\phi') = \phi$. Since $\bar{Q}$ is projective, the lifting problem

$$\begin{array}{ccc} & & \gamma_n \\ & \nearrow \bar{\phi} & \downarrow \beta_n \\ \bar{Q} & \xrightarrow{\phi'} & \text{Lie}(n) \end{array}$$

has a solution. The assertion follows by tensoring with $V^{\otimes n}$ over $R(\Sigma_n)$. $\square$

By inspecting the proof, each T_n -projective sub functor Q of L_n induces a $R(\Sigma_n)$ -projective submodule $\bar{Q}$ of $\text{Lie}(n)$. Conversely, each $R(\Sigma_n)$ -projective submodule $\bar{Q}$ of $\text{Lie}(n)$ induces a T_n -projective sub functor Q , $V \mapsto \bar{Q} \otimes_{R(\Sigma_n)} V^{\otimes n}$, of L_n . Thus we have the following:

Proposition 2.5. *There is a one-to-one correspondence between T_n -projective sub functors of L_n and $R(\Sigma_n)$ -projective submodules of $\text{Lie}(n)$.* $\square$

3. PROOF OF THEOREM 1.1

In this section, the ground field is of characteristic $p > 2$. The notation V means any fixed connected graded module such that $V_{\text{even}} = 0$ and $\dim V = p - 1$. The general graded or ungraded module is then denoted by W .

Let W be any module and let $T: W \mapsto T(W)$ be the functor from modules to Hopf algebras, where the tensor algebra $T(W)$ is Hopf by saying W primitive.

Let $B(W)$ be the sub Hopf algebra generated by $L_n(W)$ for n not a power of p . By [16, Theorem 1.5], $B(W) \subseteq B^{\max}(W)$ for any ungraded module W and so for any graded or ungraded module W . It follows that there is an epimorphism

$$(3.1) \quad q: \mathbf{k} \otimes_{B(W)} T(W) \twoheadrightarrow A^{\min}(W)$$

for any ungraded or graded module W .

We are going to determine $\mathbf{k} \otimes_{B(V)} T(V)$ and to show that the map q is in fact an isomorphism when $W = V$.

3.1. Determination of $\mathbf{k} \otimes_{B(V)} T(V)$. Let $B^{[k]}(W) = \langle B(W), L_{p^s}(W) \mid s \geq k \rangle$ be the sub Hopf algebra of $T(W)$ generated by $B(W)$ and $L_{p^s}(W)$ for $s \geq k$. Then there is a tower of sub Hopf algebras

$$\dots \subseteq B^{[k+1]}(W) \subseteq B^{[k]}(W) \subseteq \dots \subseteq B^{[1]}(W) \subseteq B^{[0]}(W) = T(W)$$

with the intersection

$$B(W) = \bigcap_{k=0}^{\infty} B^{[k]}(W).$$

Define the functor $\bar{L}_n$ by:

$$\bar{L}_n(W) = L_n(W) / \left(\sum_{i=2}^{n-2} [L_i(W), L_{n-i}(W)] \right)$$

for each $n \geq 2$. According to [16, Proposition 11.6], $Q_p B^{[1]}(W) = \bar{L}_p$. So we can identify $\bar{L}_p$ with a sub functor of L_p . Now define recursively the functor $\bar{L}_p^k$ by $\bar{L}_p^0(W) = W$, $\bar{L}_p^1(W) = \bar{L}_p(W)$ and $\bar{L}_p^{k+1}(W) = \bar{L}_p(\bar{L}_p^k(W))$. Note that $\bar{L}_p^k$ has the tensor length p^k and $\bar{L}_p^k$ can be identified with a sub functor of L_{p^k} . Observe that $(\bar{L}_p(V))_{\text{even}} = 0$ and $\dim \bar{L}_p(V) = p-1$. Thus we have the periodicity information:

$$(3.2) \quad (\bar{L}_p^k(V))_{\text{even}} = 0 \text{ and } \dim \bar{L}_p^k(V) = p-1$$

for all $k \geq 0$.

Lemma 3.1. [22, Lemma 2.37] *Let A be a connected Hopf algebra of finite type and let B be a sub Hopf algebra of A . Suppose that A is a tensor algebra as an algebra with a choice of inclusion $QA \rightarrow A$. Then there is a short exact sequence*

$$0 \longrightarrow Q(B) \longrightarrow (k \otimes_B A) \otimes Q(A) \longrightarrow I(k \otimes_B A) \longrightarrow 0.$$

Now we determine the sub Hopf algebra $B^{[k]}(V)$:

Lemma 3.2. *For each $k \geq 0$, there is a short exact sequence of Hopf algebras*

$$B^{[k+1]}(V) \hookrightarrow B^{[k]}(V) \longrightarrow E(\bar{L}_p^k(V)).$$

Moreover

- (1). $B^{[k]}(V)$ is the sub Hopf algebra of $T(V)$ generated by $\bar{L}_p^k(V)$ and $Q_j B(V)$ for $2 \leq j < p^k$.
- (2). For any possible Steenrod module structure on V , $\bar{L}_p^k(V)$ is a suspension of V .

Proof. The proof is given by induction. First consider the short exact sequence of Hopf algebra

$$B'(V) \hookrightarrow T(V) \longrightarrow \Lambda(V),$$

where $B'(V) = \mathbf{k}\square_{\Lambda(V)} T(V)$. Then $B^{[1]}(V) \subseteq B'(V)$. There is a short exact sequence

$$0 \longrightarrow Q(B'(V)) \longrightarrow \Lambda(V) \otimes V \longrightarrow I\Lambda(V) \longrightarrow 0.$$

Since $V_{\text{even}} = 0$, $E(V) = \Lambda(V)$. By the assumption of $\dim V = p-1$, we have $E(V)_j = 0$ for $j \geq p$. Thus $Q_j(B'(V)) = 0$ for $j > p$ and so $B'(V) \subseteq B^{[1]}(V)$. Hence $B'(V) = B^{[1]}(V)$. From the above exact sequence,

$$Q_p(B'(V)) \cong E_{p-1}(V) \otimes V \cong Q_p(B^{[1]}(V)) = \bar{L}_p(V)$$

is of dimension $p-1$. Since $\dim E_{p-1}(V) = 1$, $\bar{L}_p(V) = E_{p-1}(V) \otimes V$ is a suspension of V for any possible Steenrod module structure on V . Thus the assertions hold for $k=0$. Suppose that the assertion holds for k . Consider the short exact sequence of Hopf algebras

$$B''(V) \hookrightarrow B^{[k]}(V) \xrightarrow{\phi} \Lambda(\bar{L}_p^k(V)),$$

where $B''(V) = \mathbf{k}\square_{\Lambda(\bar{L}_p^k(V))} B^{[k]}(V)$. Then $B^{[k+1]}(V) \subseteq B''(V)$ because ϕ is a Hopf map which sends the generators for $B^{[k+1]}(V)$ to zero. Since $\bar{L}_p^k(V)_{\text{even}} = 0$ with $\dim \bar{L}_p^k(V) = p-1$, $\Lambda(\bar{L}_p^k(V)) = E(\bar{L}_p^k(V))$ and so there is short exact sequence

$$0 \longrightarrow Q(B''(V)) \longrightarrow E(\bar{L}_p^k(V)) \otimes Q(B^{[k]}(V)) \longrightarrow IE(\bar{L}_p^k(V)) \longrightarrow 0.$$

It follows that $Q(B''(V))_j = 0$ for $j > p^{k+1}$ and so $B''(V) \subseteq B^{[k+1]}(V)$. By the Lie action of $E(\bar{L}_p^k(V))$ on $Q(B''(V))$,

$$Q_{p^{k+1}} B^{[k+1]}(V) = Q_{p^{k+1}} B''(V) = \bar{L}_p^{k+1}(V) \cong E_{p-1}(\bar{L}_p^k(V)) \otimes \bar{L}_p^k(V).$$

Since $\dim E_{p-1}(\bar{L}_p^k(V)) = 1$, $\bar{L}_p^{k+1}(V)$ is a suspension of $\bar{L}_p^k(V)$. The induction is finished and hence the result. $\square$

Proposition 3.3. *There is a (non-functorial) coalgebra decomposition*

$$\mathbf{k} \otimes_{B(V)} T(V) \cong \bigotimes_{k=0}^{\infty} E(\bar{L}_p^k(V))$$

over any possible Steenrod algebra structure on V .

Proof. Consider the short exact sequence of coalgebras

$$(3.3) \quad \mathbf{k} \otimes_{B(V)} B^{[k+1]}(V) \xrightarrow{\quad j \quad} \mathbf{k} \otimes_{B(V)} B^{[k]}(V) \longrightarrow E(\bar{L}_p^k(V)).$$

Observe that up to tensor length $p^{k+1} - 1$

$$(\mathbf{k} \otimes_{B(V)} B^{[k]}(V))_j \cong E(\bar{L}_p^k(V))_j$$

for $j \leq p^{k+1}$. Thus there is a coalgebra cross-section

$$s: E(\bar{L}_p^k(V)) \hookrightarrow \mathbf{k} \otimes_{B(V)} B^{[k]}(V).$$

The cross-section s is a morphism over the Steenrod algebra for all possible Steenrod algebra structure on V and it is also a morphism over $\mathrm{GL}(V)$ when $\mathrm{GL}(V)$ acts on V by forgetting the grading of V .

According to [11, Theorem 1.1], $B(W)$ is a functorial coalgebra retract of $T(W)$ for any graded module W . Let $r_W: T(W) \rightarrow B(W)$ be a functorial coalgebra retraction. From $B(W) \subseteq B^{[k]}(W) \subseteq T(W)$, the restriction

$$r_V|_{B^{[k]}(W)}: B^{[k]}(V) \longrightarrow B(W)$$

is a functorial coalgebra retraction. It follows that the short exact sequence of coalgebras

$$B(W) \hookrightarrow B^{[k]}(W) \longrightarrow \mathbf{k} \otimes_{B(W)} B^{[k]}(W)$$

splits off. This gives a functorial coalgebra decomposition

$$B^{[k]}(W) \cong B(W) \otimes (\mathbf{k} \otimes_{B(W)} B^{[k]}(W))$$

for any graded module W . Thus the functor

$$W \mapsto \mathbf{k} \otimes_{B(W)} B^{[k]}(W)$$

is a functorial from modules to quasi-Hopf algebra because $\mathbf{k} \otimes_{B(W)} B^{[k]}(W)$ is a functorial coalgebra retract of the Hopf algebra functor $B^{[k]}(W)$.

Evaluating $W = V$, $\mathbf{k} \otimes_{B(V)} B^{[k]}(V)$ is a quasi-Hopf algebra over the Steenrod algebra for all possible Steenrod algebra structure on V . By [16, Lemma 5.3], the map

$$(\mathbf{k} \otimes_{B(V)} B^{[k+1]}(V)) \otimes E(\bar{L}_p^k(V)) \xrightarrow{\mu(j \otimes s)} \mathbf{k} \otimes_{B(V)} B^{[k]}(V)$$

is an isomorphism and hence the result. $\square$

3.2. Proof of Theorem 1.1. Consider the functorial short exact sequence of algebras

$$B^{\max}(W) \longrightarrow T(W) \xrightarrow{r} A^{\min}(W) = \mathbf{k} \otimes_{B^{\max}} T(W).$$

Define

$$A_n^{\min}(W) = \text{Im}(T_n(W) \hookrightarrow T(W) \xrightarrow{r} A^{\min}(W)).$$

Then

$$A^{\min}(W) = \bigoplus_{n=0}^{\infty} A_n^{\min}(W)$$

is a bigraded coalgebra and the decomposition

$$T(W) \cong B^{\max}(W) \otimes A^{\min}(W)$$

is a functorial coalgebra decomposition of bi-graded coalgebras, where the second grading on $B^{\max}(W)$ is given by $B_n^{\max}(W) = T_n(W) \cap B^{\max}(W)$. Note that

$$L_n(W) = \text{Im}(\beta_n: W^{\otimes n} \rightarrow W^{\otimes n}).$$

Denote by β_n the epimorphism $W^{\otimes n} \twoheadrightarrow L_n(W)$ if there are no confusions. The algebraic version of the James-Hopf map $H_n: T(W) \longrightarrow T(W^{\otimes n})$ is defined in [7, 16, 17].

Lemma 3.4. *There is a commutative diagram of natural transformations of functors*

$$\begin{array}{ccccccc}
 & & T & \xrightarrow{H_{p^k}} & T(T_{p^k}) & \xrightarrow{T(r)} & T(A_{p^k}^{\min}) \longrightarrow A^{\min}(A_{p^k}^{\min}) \\
 & & \uparrow & & \uparrow & & \uparrow \\
 & & \cup & & \cup & & \cup \\
 Q_{p^{k+1}}^{\max} & \hookrightarrow & L_{p^{k+1}} & \longrightarrow & L_p(T_{p^k}) & \xrightarrow{L_p(r)} & L_p(A_{p^k}^{\min}) \longrightarrow \bar{L}_p(A_{p^k}^{\min}) \\
 \parallel & & \uparrow & & \uparrow & & \uparrow \\
 Q_{p^{k+1}}^{\max} & \xrightarrow{\phi'} & T_{p^{k+1}} & \xrightarrow{\phi} & T_p(T_{p^k}) & \xrightarrow{T_p(r)} & T_p(A_{p^k}^{\min}) \\
 & & \uparrow \beta_{p^k} & & \uparrow \beta_p & & \uparrow \beta_p
 \end{array}$$

for some ϕ and ϕ' .

Proof. Since all functors in the diagram are well-defined over $\mathbb{Z}_{(p)}$, it suffices to show that the assertion holds when the ground ring is $\mathbb{Z}_{(p)}$. The top three squares commute because the maps in the top row are coalgebra maps, which send the primitives to the primitives. It is clear that the bottom right square commutes. By Proposition 2.4, there exist lifting ϕ' and ϕ such that the bottom left and middle squares commute. $\square$

Lemma 3.5. *Let $n \geq 2$ and let W be any graded module. Then the composite*

$$W^{\otimes n} = W^{\otimes 2} \otimes W^{\otimes n-2} \xrightarrow{\beta_n} L_n(W) \longrightarrow \bar{L}_n(W)$$

factors through the quotient

$$\left(W^{\otimes 2} / \langle a_1 \otimes a_2 + (-1)^{|a_1||a_2|} a_2 \otimes a_1 \rangle \right) \otimes \Lambda_{n-2}(W)$$

of $W^{\otimes n}$.

Proof. By the skew-symmetric property on the first two factors, the iterated Lie operad $\beta_n: W^{\otimes n} \rightarrow L_n(W)$ factors through the quotient

$$\left(W^{\otimes 2} / \langle a_1 \otimes a_2 + (-1)^{|a_1||a_2|} a_2 \otimes a_1 \rangle \right) \otimes W^{\otimes n-2}.$$

Consider the short exact sequence of Hopf algebras

$$B^{[1]}(W) \hookrightarrow T(W) \twoheadrightarrow \Lambda(W).$$

Note that there is a commutative diagram

$$\begin{array}{ccc} \bigoplus_{n=2}^{\infty} L_n(W) & \hookrightarrow & IB^{[1]}(W) \\ \downarrow & & \downarrow \\ \bigoplus_{n=2}^{\infty} \bar{L}_n(W) & = & QB^{[1]}(W). \end{array}$$

The assertion follows from [2, Lemma 3.12] that the Lie bracket in $T(W)$ induces an action of $\Lambda(W)$ on $QB^{[1]}(W) = \bigoplus_{n=2}^{\infty} \bar{L}_n(W)$. $\square$

Let W be a graded module such that $W_j = 0$ for $j \neq 1$. Let Σ_n act on $W^{\otimes n}$ by permuting positions in graded sense. Let $\mathrm{GL}(W_1)$ be the general linear group of the vector space W_1 . The Schur algebra [13] is defined by

$$\mathcal{S}(W) = \mathrm{Im}(\phi: \mathbf{k}(\mathrm{GL}(W)) \rightarrow \mathrm{End}(W^{\otimes n})),$$

where $\phi(f) = f^{\otimes n}$. By [5, Theorem 4.1],

$$(3.4) \quad \mathrm{Im}(\mathbf{k}(\Sigma_n) \rightarrow \mathrm{End}(W^{\otimes n})) = \mathrm{End}_{\mathcal{S}(W)}(W^{\otimes n})$$

if $\mathbf{k}$ is algebraically closed. Thus if $\mathbf{k}$ is algebraically closed, then for any $\mathbf{k}(\mathrm{GL}(W))$ -map

$$f: W^{\otimes n} \rightarrow W^{\otimes n}$$

there exists an element $\alpha \in \mathbf{k}(\Sigma_n)$ such that

$$f(a_1 \otimes \cdots \otimes a_n) = \alpha \cdot (a_1 \otimes \cdots \otimes a_n).$$

Lemma 3.6. *Let $\bar{V}$ be a graded module such that $\dim \bar{V}_1 = p-1$ and $\bar{V}_j = 0$ for $j \neq 1$. Let $\mathrm{GL}(\bar{V})$ be the general linear group with the action on $\bar{V}$. Assume that the ground field $\mathbf{k}$ is algebraically closed. Then, as a morphism over $\mathbf{k}(\mathrm{GL}(\bar{V}))$, the composite*

$$q: T_p(\bar{V}) = \bar{V}^{\otimes p} \xrightarrow{\beta_p} L_p(\bar{V}) \twoheadrightarrow \bar{L}_p(\bar{V}) \xrightarrow{\cong} \bar{V} \otimes E_{p-1}(\bar{V})$$

does NOT have a cross-section.

Proof. Suppose that there exists a $\mathbf{k}(\mathrm{GL}(\bar{V}))$ -map $\phi: \bar{V} \otimes E_{p-1}(\bar{V}) \rightarrow \bar{V}^{\otimes p}$ such that the composite $q \circ \phi$ is the identity map of $\bar{V} \otimes E_{p-1}(\bar{V})$. We are going to find a contradiction.

Let $\{x_1, x_2, \dots, x_{p-1}\}$ be a basis for $\bar{V}$. Note that

$$\dim E_{p-1}(\bar{V}) = 1$$

with a basis $\{(x_1 \wedge x_2 \wedge \cdots \wedge x_{p-1})\}$. The set

$$\{x_i \otimes (x_1 \wedge x_2 \wedge \cdots \wedge x_{p-1}) \mid 1 \leq i \leq p-1\}$$

forms a basis for $\bar{V} \otimes E_{p-1}(\bar{V})$. Since the composite

$$\bar{V}^{\otimes p} \xrightarrow{q} \bar{V} \otimes E_{p-1}(\bar{V}) \xrightarrow{\phi} \bar{V}^{\otimes p}$$

is an endomorphism over $\mathbf{k}(\mathrm{GL}(\bar{V}))$, from equation (3.4) there exists an element $\alpha \in \mathbf{k}(\Sigma_p)$ such that

$$(3.5) \quad \phi \circ q(w) = \alpha \cdot w$$

for $w \in \bar{V}^{\otimes p}$. By counting the occurrence of x_j 's, $\phi(x_i \otimes (x_1 \wedge \cdots \wedge x_{p-1}))$ is a linear combination of monomials $x_{i_1} x_{i_2} \cdots x_{i_p}$ in which x_i occurs twice and x_j occurs once for each $j \neq i$. By Lemma 3.5, there is a commutative diagram

$$\begin{array}{ccc} \bar{V}^{\otimes p} & \xrightarrow{\beta_p} & L_p(\bar{V}) \\ \downarrow \pi & & \downarrow \bar{\pi} \\ (\bar{V}^{\otimes 2} / \langle x_i x_j - x_j x_i \rangle) \otimes E_{p-2}(\bar{V}) & \xrightarrow{g} & \bar{L}_p(\bar{V}) = \bar{V} \otimes E_{p-1}(\bar{V}). \end{array}$$

By counting the occurrence of x_j 's,

$$(3.6) \quad \begin{aligned} & \pi \circ \phi(x_1 \otimes (x_1 \wedge x_2 \wedge \cdots \wedge x_{p-1})) \\ &= k_1 x_1^2 \otimes (x_2 \wedge x_3 \wedge \cdots \wedge x_{p-1}) \\ &+ \sum_{i=2}^{p-1} k_i x_1 x_i \otimes (x_1 \wedge \cdots \wedge x_{i-1} \wedge x_{i+1} \wedge \cdots \wedge x_{p-1}). \end{aligned}$$

for some $k_i \in \mathbf{k}$. Note that $x_i \wedge x_j = -x_j \wedge x_i$. By interchanging x_i and x_{i+1} , there is an equation

$$k_i = -k_{i+1}$$

for $i \geq 2$. By equation (3.5), the map

$$\pi \circ \phi: \bar{V} \otimes E_{p-1}(\bar{V}) \longrightarrow (\bar{V}^{\otimes 2} / \langle x_i x_j - x_j x_i \rangle) \otimes E_{p-2}(\bar{V})$$

is a morphism over the Steenrod algebra by regarding $\bar{V}$ as *any graded module over the Steenrod algebra* with $\bar{V}_{\text{even}} = 0$. Take the Steenrod module structure on $\bar{V}$ by setting $P_*^1 x_1 = x_2$, $P_*^i x_1 = 0$ for $i > 1$ and $P_*^k x_j = 0$ for $k \geq 1$ and $j > 1$. Then

$$\begin{aligned} & P_*^2 x_1 x_i \otimes (x_1 \wedge \cdots \wedge x_{i-1} \wedge x_{i+1} \wedge \cdots \wedge x_{p-1}) \\ &= x_2 x_i \otimes (x_2 \wedge \cdots \wedge x_{i-1} \wedge x_{i+1} \wedge \cdots \wedge x_{p-1}) \\ &= 0 \end{aligned}$$

for $i \geq 3$. Note that

$$P_*^2 x_1 \otimes (x_1 \wedge x_2 \wedge \cdots \wedge x_{p-1}) = x_2 \otimes (x_2 \wedge x_2 \wedge \cdots \wedge x_{p-1}) = 0.$$

By applying P_*^2 to equation (3.6),

$$\begin{aligned} 0 &= P_*^2(k_1 x_1^2 \otimes (x_2 \wedge x_3 \wedge \cdots \wedge x_{p-1}) + k_2 x_1 x_2 \otimes (x_1 \wedge x_3 \wedge \cdots \wedge x_{p-1})) \\ &= k_1 x_1^2 \otimes (x_2 \wedge x_3 \wedge \cdots \wedge x_{p-1}) + k_2 x_2^2 \otimes (x_2 \wedge x_3 \wedge \cdots \wedge x_{p-1}). \end{aligned}$$

Thus

$$k_1 = -k_2$$

and so

$$\pi \circ \phi(x_1 \otimes (x_1 \wedge x_2 \wedge \cdots \wedge x_{p-1})) = k_1 \sum_{i=1}^{p-1} (-1)^{i-1} x_1 x_i \otimes (x_1 \wedge \cdots \wedge x_{i-1} \wedge x_{i+1} \wedge \cdots \wedge x_{p-1}).$$

Define

$$\alpha_1 = x_1 \otimes x_1 \otimes x_2 \otimes x_3 \otimes \cdots \otimes x_{p-1} \in \bar{V}^{\otimes p}$$

$$\alpha_i = x_1 \otimes x_i \otimes x_1 \otimes \cdots \otimes x_{i-1} \otimes x_{i+1} \otimes \cdots \otimes x_{p-1} \in \bar{V}^{\otimes p}$$

for $2 \leq i \leq p-1$. Then

$$\begin{aligned} \bar{\pi}\beta_p(\alpha_1) &= \bar{\pi}([x_1, x_1], \dots, x_{p-1}) \\ &= 2x_1 \otimes (x_1 \wedge x_2 \wedge x_3 \wedge \cdots \wedge x_{p-1}). \end{aligned}$$

For $i \geq 2$,

$$\begin{aligned} \bar{\pi}\beta_p(\alpha_i) &= \bar{\pi}([x_1, x_i], x_1, \dots, x_{i-1}, x_{i+1}, x_{p-1}) \\ &= (-1)^{i-1} x_1 \otimes (x_1 \wedge x_2 \wedge x_3 \wedge \cdots \wedge x_{p-1}). \end{aligned}$$

It follows that

$$\begin{aligned} &q \circ \phi(x_1 \otimes (x_1 \wedge x_2 \wedge \cdots \wedge x_{p-1})) \\ &= g \circ \pi \circ \phi(x_1 \otimes (x_1 \wedge x_2 \wedge \cdots \wedge x_{p-1})) \\ &= k_1 g \left(\sum_{i=1}^{p-1} (-1)^{i-1} x_1 x_i \otimes (x_1 \wedge \cdots \wedge x_{i-1} \wedge x_{i+1} \wedge \cdots \wedge x_{p-1}) \right) \\ &= k_1 g \left(\sum_{i=1}^{p-1} (-1)^{i-1} \pi(\alpha_i) \right) \\ &= k_1 \sum_{i=1}^{p-1} (-1)^{i-1} g \circ \pi(\alpha_i) \\ &= k_1 (\bar{\pi}\beta_p(\alpha_1) + \sum_{i=2}^{p-1} (-1)^{i-1} \bar{\pi}\beta_p(\alpha_i)) \\ &= pk_1 x_1 \otimes (x_1 \wedge x_2 \wedge x_3 \wedge \cdots \wedge x_{p-1}) \\ &= 0, \end{aligned}$$

which contradicts to the assumption that $q \circ \phi$ is the identity map. This finishes the proof. $\square$

Proof of Theorem 1.1. By Proposition 3.3, it suffices to show that the epimorphism

$$\mathbf{k} \otimes_{B(V)} T(V) \longrightarrow A^{\min}(V)$$

is an isomorphism.

Since the functor $A^{\min}$ only depends on the characteristic of the ground field, we may assume that the ground field $\mathbf{k}$ is algebraically closed. Denote by ϕ_k the epimorphism $\mathbf{k} \otimes_{B(V)} T(V) \longrightarrow A^{\min}(V)$. By Proposition 3.3,

$$\bar{L}_p^k(V) = P_{p^k}(\mathbf{k} \otimes_{B(V)} T(V))$$

for each $k \geq 0$. By equation (3.1), $A^{\min}(V)$ is functorial coalgebra retract of $\mathbf{k} \otimes_{B(V)} T(V)$. It suffices to show by induction that the retraction map

$$\phi_k|_{\bar{L}_p^k(V)}: \bar{L}_p^k(V) = P_{p^k}(\mathbf{k} \otimes_{B(V)} T(V)) \rightarrow P_{p^k} A^{\min}(V)$$

is an isomorphism for each $k \geq 0$. The statement holds for $k = 0$. Suppose that the statement holds for $s \leq k$ with $k > 0$ and consider

$$\phi_{k+1}|_{\bar{L}_p^{k+1}(V)}: \bar{L}_p^{k+1}(V) = P_{p^{k+1}}(\mathbf{k} \otimes_{B(V)} T(V)) \rightarrow P_{p^{k+1}} A^{\min}(V).$$

By choosing V to a module over the Steenrod algebra with a basis given by $\{v, P_*^1 v, P_*^2 v, \dots, P_*^{p-2} v\}$, then V is an indecomposable module over the Steenrod algebra. By Lemma 3.2, $\bar{L}_p^{k+1}(V)$ is a suspension of V as a module over the Steenrod algebra. Thus $\bar{L}_p^{k+1}(V)$ is an indecomposable module over the Steenrod algebra. It follows that the retraction

$$\phi_{k+1}|_{\bar{L}_p^{k+1}(V)}: \bar{L}_p^{k+1}(V) = P_{p^{k+1}}(\mathbf{k} \otimes_{B(V)} T(V)) \rightarrow P_{p^{k+1}} A^{\min}(V)$$

is either identically zero or an isomorphism.

Suppose that $\phi_{k+1}|_{\bar{L}_p^{k+1}(V)} = 0$. Then

$$P_{p^{k+1}}A^{\min}(V) = 0$$

and so

$$L_{p^{k+1}}(V) \subseteq B^{\max}(V).$$

By Lemma 3.2, $B^{[k+1]}(V)$ is the sub Hopf algebra of $T(V)$ generated by $\bar{L}_p^{k+1}(V)$ and $L_n(V)$ for n not a power of p . It follows that

$$(3.7) \quad B^{[k+1]}(V) \subseteq B^{\max}(V)$$

and so

$$PA^{\min}(V) \subseteq \bigoplus_{s=1}^k \bar{L}_p^s(V).$$

From the induction. $\bar{L}_p^s(V) \subseteq A^{\min}(V)$ for each $s \leq k$. Thus

$$PA^{\min}(V) = \bigoplus_{s=1}^k \bar{L}_p^s(V).$$

It follows that there is a (*non-functorial*) isomorphism of coalgebra

$$A^{\min}(V) \cong \bigotimes_{s=0}^k E(\bar{L}_p^s(V)).$$

By computing Poincaré series, the inequality (3.7) becomes the equality

$$B^{[k+1]}(V) = B^{\max}(V).$$

Thus

$$Q_{p^{k+1}}^{\max}(V) = Q_{p^{k+1}}B^{\max}(V) = Q_{p^{k+1}}B^{[k+1]}(V) \cong \bar{L}_p^{k+1}(V).$$

Observe that the inclusion

$$\bar{L}_p^k(V) = P_{p^k}A^{\min}(V) \hookrightarrow A_{p^k}^{\min}(V)$$

is an isomorphism and $\bar{L}_p^{k+1}(V) = \bar{L}_p(\bar{L}_p^k(V))$. The composite of natural transformations

$$(3.8) \quad Q_{p^{k+1}}^{\max} \hookrightarrow L_{p^{k+1}} \longrightarrow L_p(T_{p^k}) \xrightarrow{L_p(r)} L_p(A_{p^k}^{\min}) \longrightarrow \bar{L}_p(A_{p^k}^{\min})$$

becomes an isomorphism by evaluating on the graded module V . Write $\bar{V}$ for $\bar{L}_p^k(V) = P_{p^k}A^{\min}(V) = A_{p^k}^{\min}(V)$. By Lemma 3.4, the composite of the first three natural transformations in equation (3.8) admits a lifting of natural transformation into $T_p(A_{p^k}^{\min})$ via the epimorphism $\beta_p: T_p(A_{p^k}^{\min}) \longrightarrow L_p(A_{p^k}^{\min})$ and so the composite

$$q: T_p(\bar{V}) = \bar{V}^{\otimes p} \xrightarrow{\beta_p} L_p(\bar{V}) \longrightarrow \bar{L}_p(\bar{V}) \xrightarrow{\cong} \bar{V} \otimes E_{p-1}(\bar{V})$$

admits a cross-section over the general linear group algebra $\mathbf{k}(\mathrm{GL}(\bar{V}))$ by forgetting the grading of $\bar{V}$. This is impossible according to Lemma 3.6.

Thus $\phi_{k+1}|_{\bar{L}_p^{k+1}(V)} \neq 0$ and so $\phi_{k+1}|_{\bar{L}_p^{k+1}(V)}$ must be an isomorphism. The induction is finished and hence the result. $\square$

4. THE GEOMETRY OF NATURAL COALGEBRA DECOMPOSITIONS

4.1. Geometric Realizations. Denote by $\mathbf{CoH}$ the category of p -local simply connected co- H -spaces of finite type and co- H -maps. Let $[\Omega, \Omega]_{\mathbf{CoH}}$ be the group of natural transformations of the functor Ω from $\mathbf{CoH}$ to the homotopy category of spaces.

Theorem 4.1 (Geometric Realization Theorem). [15] *Let Y be any simply connected co- H -space of finite type and let*

$$T(V) \cong A(V) \otimes B(V)$$

any natural coalgebra decomposition for ungraded modules over $\mathbb{Z}/p$. Then there exist homotopy functors $\bar{A}$ and $\bar{B}$ from $\mathbf{CoH}$ to spaces such that

- (1). *there is a functorial decomposition*

$$\Omega Y \simeq \bar{A}(Y) \times \bar{B}(Y)$$

- (2). *On mod p homology the decomposition*

$$H_*(\Omega Y) \cong H_*(\bar{A}(Y)) \otimes H_*(\bar{B}(Y))$$

is with respect to the augmentation ideal filtration

- (3). *On mod p homology*

$$E^0 H_*(\bar{A}(Y)) = A(\Sigma^{-1} \bar{H}_*(Y)) \text{ and } E^0 H_*(\bar{B}(Y)) = B(\Sigma^{-1} \bar{H}_*(Y)),$$

where A and B are the canonical extensions of the functors A and B for graded modules $\square$

For the functors $A^{\min}$ and $B^{\max}$, we have the following geometric realization theorem.

Theorem 4.2. [15] *There exist homotopy functors $\bar{A}^{\min}$, $\bar{Q}_n^{\max}$, $n \geq 2$, from $\mathbf{CoH}$ to spaces such that for any p -local simply connected co- H space Y of finite type the following hold:*

- 1) $\bar{B}^{\max}(Y) = \Omega(\bigvee_{n=2}^{\infty} \bar{Q}_n^{\max}(Y))$.
- 2) *There is a functorial decomposition*

$$\Omega Y \simeq \bar{A}^{\min}(Y) \times \bar{B}^{\max}(Y).$$

- 3) *On mod p homology the decomposition*

$$H_*(\Omega Y) \cong H_*(\bar{A}^{\min}(Y)) \otimes H_*(\bar{B}^{\max}(Y))$$

is with respect to the augmentation ideal filtration.

- 4) *On mod p homology*

$$\begin{aligned} E^0 H_*(\bar{A}^{\min}(Y)) &= A^{\min}(\Sigma^{-1} \bar{H}_*(Y)), \\ E^0 H_*(\bar{B}^{\max}(Y)) &= B^{\max}(\Sigma^{-1} \bar{H}_*(Y)), \\ E^0 H_*(\bar{Q}^{\max}(Y)) &= Q^{\max}(\Sigma^{-1} \bar{H}_*(Y)). \end{aligned}$$

$\square$

4.2. Suspension Splitting Theorems. In this subsection, we review the suspension splitting theorems.

A *graded space* means a space W with a homotopy decomposition

$$\phi_W: W \xrightarrow{\simeq} \bigvee_{n=1}^{\infty} W_n.$$

For any graded space W , the homology $\bar{H}_*(W)$ is filtered by

$$I^t \bar{H}_*(W) = \phi_*^{-1}(\bar{H}_*(\bigvee_{n=t}^{\infty} W_n))$$

for $t \geq 1$. A *graded co- H space* means a graded space W such that W is a co- H space. Thus each factor W_n is also a co- H space. The following lemma gives a general criterion for decomposing the retracts of graded co- H spaces in term of grading factors.

Lemma 4.3. [6] *Let W be a simply connected p -local graded co- H space of finite type. Let $f: W \rightarrow W$ be a self-map such that on mod p homology*

- 1) $f_*: \bar{H}_*(W) \rightarrow \bar{H}_*(W)$ *preserves the filtration.*
- 2) *The induced bigraded map $E^0 f_*$ is an idempotent:*

$$E^0 f_* \circ E^0 f_* = E^0 f_*: E^0 H_*(W) \rightarrow E^0 H_*(W).$$

Let $A(f) = \text{hocolim}_f W$ be the homotopy colimit and let $A_n(f) = \text{hocolim}_{g_n} W_n$, where g_n is the composite

$$g_n: W_n \hookrightarrow \bigvee_{k=1}^{\infty} W_k \xrightarrow{\phi_W^{-1}} W \xrightarrow{f} W \xrightarrow{\phi_W} \bigvee_{k=1}^{\infty} W_k \longrightarrow W_n.$$

Then there is a canonical homotopy decomposition of the homotopy colimit

$$A(f) \simeq \bigvee_{n=1}^{\infty} A_n(f)$$

such that

$$\bar{H}_*(A_n(f)) \cong \text{Im}(E_n^0 f_*: E^0 H_*(W) \cong \bar{H}_*(W_n) \rightarrow E^0 H_*(W) \cong \bar{H}_*(W_n)).$$

□

Let X be any path-connected space. Let $H_*(\Omega \Sigma X)$ be filtered by the powers of the augmentation ideal filtration. From the classical suspension splitting Theorem [10]

$$\phi: \Sigma \Omega \Sigma X \xrightarrow{\simeq} \bigvee_{n=1}^{\infty} \Sigma X^{(n)},$$

$\Sigma \Omega \Sigma X$ is a simply graded co- H space and the filtration

$$I^t \bar{H}_*(\Sigma \Omega \Sigma X) = \phi_*^{-1}(\bar{H}_*(\bigvee_{n=t}^{\infty} \Sigma X^{(n)}))$$

coincide with (the suspension) of the augmentation ideal filtration of $H_*(\Omega \Sigma X)$.

Let W and W' be graded spaces. Then $W \wedge W'$ is a graded space with homotopy equivalence

$$W \wedge W' \xrightarrow[\simeq]{\phi_W \wedge \phi_{W'}} \left(\bigvee_{n=1}^{\infty} W_n \right) \wedge \left(\bigvee_{n=1}^{\infty} W'_n \right) = \bigvee_{n=1}^{\infty} \bigvee_{i=1}^{n-1} W_i \wedge W'_{n-i}$$

and the homology $\bar{H}_*(W \wedge W') = \bar{H}_*(W) \otimes \bar{H}_*(W')$ is the tensor product of filtered modules. Moreover if W is a graded co- H space, then $W \wedge W'$ is a graded co- H space.

Recall that [16] any natural coalgebra retract $A(V)$ of $T(V)$ for ungraded modules admits the tensor length decomposition

$$A(W) = \bigoplus_{n=0}^{\infty} A_n(W)$$

for any graded or ungraded module W .

Theorem 4.4 (Suspension Splitting Theorem [6]). *Let $A(V)$ be any natural coalgebra retract of $T(V)$ for any ungraded modules V and let $\bar{A}$ be the geometric realization of A . Then for any p -local simply connected co- H space Y of finite type and any p -local path-connected co- H -space Z , there is a functorial splitting*

$$Z \wedge \bar{A}(Y) \simeq \bigvee_{n=1}^{\infty} [Z \wedge \bar{A}(Y)]_n$$

such that

$$\bar{H}_*([Z \wedge \bar{A}(Y)]_n) \cong \bar{H}_*(Z) \otimes A_n(\Sigma^{-1} \bar{H}_*(Y))$$

for each $n \geq 1$. In particular, for a p -local simply connected co- H -space Y there is a functorial suspension splitting

$$\Sigma \bar{A}(Y) \simeq \bigvee_{n=1}^{\infty} \bar{A}_n(Y)$$

such that

$$\Sigma^{-1} \bar{H}_*(\bar{A}_n(Y)) \cong A_n(\Sigma^{-1} \bar{H}_*(Y))$$

for each $n \geq 1$. □

4.3. Hopf Invariants. In this section, we review the results in Hopf invariants from [6]. Let $A(V)$ be a natural coalgebra retract of $T(V)$ for ungraded modules. From the suspension splitting theorem (Theorem 4.4), there is a decomposition

$$\Sigma \bar{A}(Y) \simeq \bigvee_{n=1}^{\infty} \bar{A}_n(Y)$$

and so it induces Hopf invariants

$$H_n: \bar{A}(Y) \longrightarrow \Omega(\bar{A}_n(Y)).$$

For computational purpose on homology, it is useful to make a particular choice of Hopf invariants H_n . Let

$$\mathcal{H}: \Sigma \Omega \Sigma X \simeq \Sigma J(X) \xrightarrow{\simeq} \bigvee_{n=1}^{\infty} \Sigma X^{(n)}$$

be the fat James-Hopf invariants. Let $H_*(\Omega \Sigma X)$ be filtered by the products of the augmentation ideal and let $H_*(\bigvee_{n=1}^{\infty} X^{(n)})$ be filtered by

$$\bigoplus_{t \geq n} H_*(X^{(t)}).$$

By [17, Proposition 3.7], the isomorphism

$$\mathcal{H}_* : H_*(\Sigma\Omega\Sigma X) \longrightarrow H_*\left(\bigvee_{n=1}^{\infty} \Sigma X^{(n)}\right)$$

preserves the filtration. Note that the composite

$$\Sigma\Omega\Sigma X \xrightarrow{\mathcal{H}} \bigvee_{n=1}^{\infty} \Sigma X^{(n)} \xrightarrow{\text{proj}} \Sigma X^{(n)}$$

is the James-Hopf invariant H_n . Let $Y \in \mathbf{CoH}$ be a simply connected p -local co- H -space of finite type with the cross-section map $s_{\mu'} : Y \longrightarrow \Sigma\Omega Y$. Let f be the composite

$$\Sigma\Omega\Sigma\Omega Y \xrightarrow{\Sigma\Omega\sigma} \Sigma\Omega Y \xrightarrow{\Sigma\Omega s_{\mu'}} \Sigma\Omega\Sigma\Omega Y.$$

By Lemma 4.3, the space $[\Sigma\Omega Y]_n$ is defined to be the homotopy colimit of the self map $g_n(Y)$ given by the composite

$$\begin{array}{ccc} \Sigma(\Omega Y)^{(n)} & \xrightarrow{\quad \subset \quad} & \bigvee_{k=1}^{\infty} \Sigma(\Omega Y)^{(k)} \\ \xrightarrow{\Sigma\Omega(s_{\mu'} \circ \sigma)} & & \xrightarrow{\mathcal{H}^{-1}} \Sigma\Omega\Sigma\Omega Y \\ \xrightarrow{\text{proj}} & \Sigma\Omega\Sigma\Omega Y & \xrightarrow{\mathcal{H}} \bigvee_{k=1}^{\infty} \Sigma(\Omega Y)^{(k)} \\ & \Sigma(\Omega Y)^{(n)}. & \end{array}$$

Proposition 4.5. *For any $Y \in \mathbf{CoH}$, the composite*

$$\begin{array}{ccc} \mathcal{H}^Y : \Sigma\Omega Y & \xrightarrow{\Sigma\Omega s_{\mu'}} & \Sigma\Omega\Sigma\Omega Y \\ \longrightarrow & \bigvee_{n=1}^{\infty} \text{hocolim}_{g_n} \Sigma(\Omega Y)^{(n)} & \xrightarrow[\cong]{\mathcal{H}} \bigvee_{n=1}^{\infty} \Sigma(\Omega Y)^{(n)} \end{array}$$

is a homotopy equivalence. $\square$

Now define the n th Hopf invariant H_n^Y to be the adjoint map to the composite

$$(4.1) \quad \Sigma\Omega Y \xrightarrow{\mathcal{H}^Y} \bigvee_{n=1}^{\infty} \text{hocolim}_{g_n} \Sigma(\Omega Y)^{(n)} \xrightarrow{\text{proj}} \text{hocolim}_{g_n} \Sigma(\Omega Y)^{(n)} = [\Sigma\Omega Y]_n.$$

Theorem 4.6. [6] *For any $Y \in \mathbf{CoH}$, there is a commutative diagram*

$$\begin{array}{ccc} E^0 H_*(\Omega Y) & \xrightarrow{E^0 H_{n*}^Y} & E^0 H_*(\Omega[\Sigma\Omega Y]_n) \\ \uparrow \cong & & \uparrow \cong \\ T(\Sigma^{-1} \bar{H}_*(Y)) & \xrightarrow{H_n} & T((\Sigma^{-1} \bar{H}_*(Y))^{\otimes n}), \end{array}$$

where $H_n : T(V) \rightarrow T(V^{\otimes n})$ is the algebraic James-Hopf map. $\square$

5. PROOF OF THEOREM 1.5

5.1. Proof of Theorem 1.5. Now we give the proof of Theorem 1.5. All spaces are localized at $p > 2$. For a natural coalgebra retract A of T , let $\bar{A}$ be its geometric realization. According to the suspension splitting theorem 4.4, there is a decomposition

$$\Sigma\bar{A}(Y) \simeq \bigvee_{n=1}^{\infty} \bar{A}_n(Y).$$

In particular, we have notations $\bar{A}^{\min}$ and $\bar{A}_n^{\min}$. Let X be a path-connected finite complex. Define

$$b_X = \sum_{q=1}^{\infty} q \dim \bar{H}_q(X; \mathbb{Z}/p).$$

Lemma 5.1. *Let Y be a simply connected co- H -space such that $\bar{H}_{\text{odd}}(Y) = 0$ and $\dim \bar{H}_*(Y) = p - 1$. Then*

$$\bar{A}_{p^k-1}^{\min}(Y) \simeq S^{(b_Y - p + 1)\frac{p^k-1}{p-1} + 1}.$$

Proof. Let $V = \Sigma^{-1} \bar{H}_*(Y)$. Then $V_{\text{even}} = 0$ and $\dim V = p - 1$. By Theorem 1.1,

$$A^{\min}(V) = \bigotimes_{k=0}^{\infty} E(\bar{L}_p^k(V)).$$

By considering tensor length $p^k - 1$, we have

$$A_{p^k-1}^{\min}(V) = E_{p-1}(V) \otimes E_{p-1}(\bar{L}_p(V)) \otimes \cdots \otimes E_{p-1}(\bar{L}_p^{k-1}(V)),$$

which is a one-dimensional module. The assertion follows from Theorem 4.4. $\square$

Let $\bar{Q}_n^{\max}$ the geometric realization of the indecomposables of $B^{\max}(V)$ as given in Theorem 4.2. Note that Y is a retract of $\Sigma \bar{A}^{\min}(Y)$ by Theorem 4.4.

Lemma 5.2. *Let Y be any p -local simply connected co- H space of finite type. Then*

(1). *there is a splitting cofibre sequence*

$$\bar{Q}^{\max}(Y) \longrightarrow \bar{A}^{\min}(Y) \wedge Y \longrightarrow (\Sigma \bar{A}^{\min}(Y))/Y.$$

Thus there is a decomposition $Y \wedge \bar{A}^{\min}(Y) \simeq \bar{Q}^{\max}(Y) \vee (\Sigma \bar{A}^{\min}(Y))/Y$.

(2). *There is a decomposition*

$$[Y \wedge \bar{A}^{\min}(Y)]_n \simeq \bar{Q}_n^{\max}(Y) \vee \bar{A}_n^{\min}(Y).$$

(3). *If $Y = \Sigma X$, then*

$$[Y \wedge \bar{A}^{\min}(Y)]_n \simeq X \wedge \bar{A}_{n-1}^{\min}(Y).$$

(4). *If $\bar{A}_{n-1}^{\min}(Y) \simeq \Sigma Z$ for some Z , then*

$$[Y \wedge \bar{A}^{\min}(Y)]_n \simeq Y \wedge Z.$$

(5). $\bar{A}_p^{\min}(Y) \simeq [Y \wedge \bar{A}^{\min}(Y)]_p$.

Proof. (1) and (2). Since $\bar{B}^{\max}(Y) \simeq \Omega \bar{Q}^{\max}(Y)$, there is a fibre sequence

$$\Omega Y \xrightarrow{\partial} \bar{A}^{\min}(Y) \xrightarrow{\simeq *} \bar{Q}^{\max}(Y) \longrightarrow Y.$$

There is a (right) action of ΩY on $\bar{A}^{\min}(Y)$ such that the diagram

$$\begin{array}{ccc} \Omega Y \times \Omega Y & \longrightarrow & \Omega Y \\ \downarrow \partial \times \text{id}_{\Omega Y} & & \downarrow \partial \\ \bar{A}^{\min}(Y) \times \Omega Y & \longrightarrow & \bar{A}^{\min}(Y) \end{array}$$

commutes. Consider the commutative diagram

$$\begin{array}{ccccccc}
 & & & & \Sigma\Omega\bar{Q}^{\max}(Y) & \longleftarrow \bar{Q}^{\max}(Y) & \\
 & & & & \downarrow & & \\
 (\Omega Y) \wedge Y & \xrightarrow{\text{id}_{\Omega Y} \wedge s} & \Sigma\Omega Y \wedge \Omega Y & \xrightarrow{H} & \Sigma\Omega Y & \longrightarrow & (\Sigma\Omega Y)/Y \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \bar{A}^{\min}(Y) \wedge Y & \xrightarrow{\text{id}_{\bar{A}^{\min}(Y)} \wedge s} & \bar{A}^{\min}(Y) \wedge \Sigma\Omega Y & \xrightarrow{H} & \Sigma\bar{A}^{\min}(Y) & \longrightarrow & (\Sigma\bar{A}^{\min}(Y))/Y,
 \end{array}$$

where H is the Hopf construction. Observe that the composite of the middle row is a homotopy equivalence. It follows that there is a sequence

$$(5.1) \quad \bar{Q}^{\max}(Y) \xrightarrow{f} \bar{A}^{\min}(Y) \wedge Y \xrightarrow{g} (\Sigma\bar{A}^{\min}(Y))/Y$$

such that $g \circ f \simeq *$, where g is the composite of the bottom rows and f is the composite of the maps from top row down to the middle right composing with the homotopy inverse from $(\Sigma\Omega Y)/Y$ to $\Omega Y \wedge Y$ and the left column map. Let C_f be the homotopy cofibre of the map f . By Lemma 3.1, the resulting map

$$C_f \longrightarrow (\Sigma\bar{A}^{\min}(Y))/Y$$

is a homotopy equivalence. Thus Sequence (5.1) is a cofibre sequence. Let r be the composite

$$(5.2) \quad r: \bar{A}^{\min}(Y) \wedge Y \xrightarrow{s^{\min} \wedge \text{id}_Y} (\Omega Y) \wedge Y \xrightarrow{H \circ (\text{id}_{\Omega Y} \wedge s)} \Sigma\Omega Y \longrightarrow \Sigma\Omega\bar{Q}^{\max}(Y) \xrightarrow{\sigma} \bar{Q}^{\max}(Y).$$

By the proof of [22, Lemma 2.37], on homology

$$(r \circ f)_*: \bar{H}_*(\bar{Q}^{\max}(Y)) \longrightarrow \bar{H}_*(\bar{Q}^{\max}(Y))$$

is an isomorphism. Thus $f: \bar{Q}^{\max}(Y) \rightarrow \bar{A}^{\min}(Y) \wedge Y$ admits a retraction and hence assertion (1). Let $H_*(\Omega Y)$ be filtered by the products of the augmentation ideal. Then $H_*(\bar{A}^{\min}(Y))$ and $H_*(\bar{Q}^{\max}(Y))$ have the induced filtration. Observe that the maps f, g and r induces filtration preserving maps on homology. Thus the decomposition

$$Y \wedge \bar{A}^{\min}(Y) \simeq \bar{Q}^{\max}(Y) \vee (\Sigma\bar{A}^{\min}(Y))/Y$$

induces a graded decomposition and hence assertion (2).

(3) By Theorem 4.4,

$$\Sigma\bar{A}^{\min}(Y) \simeq \bigvee_{k=1}^{\infty} \bar{A}_k^{\min}(Y).$$

Let $s: \bar{A}_{n-1}^{\min}(Y) \hookrightarrow \Sigma\bar{A}^{\min}(Y)$ be the inclusion. Then composite

$$\begin{aligned}
 X \wedge \bar{A}_{n-1}^{\min}(Y) & \xrightarrow{\text{id}_X \wedge s} X \wedge \Sigma\bar{A}^{\min}(Y) \\
 & \simeq Y \wedge \bar{A}^{\min}(Y) \\
 & \simeq \bigvee_{k=2}^{\infty} [Y \wedge \bar{A}_k^{\min}(Y)]_k \\
 & \xrightarrow{\text{proj.}} [Y \wedge \bar{A}^{\min}(Y)]_n
 \end{aligned}$$

induces an isomorphism on homology and hence assertion (3).

(4) Let $s: Y \rightarrow \Sigma\Omega Y$ be a cross-section to the evaluation map $\sigma: \Sigma\Omega Y \rightarrow Y$. Let f be the idempotent

$$\Sigma\Omega Y \wedge \bar{A}^{\min}(Y) \xrightarrow{\sigma \wedge \text{id}_{\bar{A}^{\min}(Y)}} Y \wedge \bar{A}^{\min}(Y) \xrightarrow{s \wedge \text{id}_{\bar{A}^{\min}(Y)}} \Sigma\Omega Y \wedge \bar{A}^{\min}(Y)$$

and let g_n be the composite

$$\begin{array}{ccc} \Omega Y \wedge \bar{A}_{n-1}^{\min}(Y) & \simeq & [\Sigma\Omega Y \wedge \bar{A}^{\min}(Y)]_n \\ & \hookrightarrow & \Sigma\Omega Y \wedge \bar{A}^{\min}(Y) \\ & \xrightarrow{f} & \Sigma\Omega Y \wedge \bar{A}^{\min}(Y) \\ & \simeq & \Omega Y \wedge \Sigma \bar{A}^{\min}(Y) \\ & \xrightarrow{\text{proj.}} & \Omega Y \wedge \bar{A}_{n-1}^{\min}(Y), \end{array}$$

where the top homotopy equivalence follows from assertion (3) with $X = \Omega Y$. According to Lemma 4.3,

$$[Y \wedge \bar{A}^{\min}(Y)]_n \simeq \text{hocolim}_{g_n} \Omega Y \wedge \bar{A}_{n-1}^{\min}(Y)$$

with the canonical retraction

$$r: \Omega Y \wedge \bar{A}_{n-1}^{\min}(Y) \longrightarrow [Y \wedge \bar{A}^{\min}(Y)]_n.$$

Now the composite

$$Y \wedge Z \xrightarrow{s \wedge \text{id}_Z} \Sigma\Omega Y \wedge Z \simeq \Omega Y \wedge \bar{A}_{n-1}^{\min}(Y) \xrightarrow{r} [Y \wedge \bar{A}^{\min}(Y)]_n$$

induces an isomorphism on homology and so it is a homotopy equivalence. Assertion (4) follows.

(5). According to [16, Section 11.2, p.97], $Q_p^{\max}(V) = 0$. Thus

$$\bar{H}_*(\bar{Q}_p^{\max}(Y)) = 0$$

and so $\bar{Q}_p^{\max}(Y) \simeq *$. By assertion (2),

$$\bar{A}_p^{\min}(Y) \simeq [Y \wedge \bar{A}^{\min}(Y)]_p$$

and hence the result. $\square$

Lemma 5.3. *Let Y be a simply connected co- H -space such that $\bar{H}_{\text{odd}}(Y) = 0$ and $\dim \bar{H}_*(Y) = p - 1$. Then*

$$\bar{A}_p^{\min}(Y) \simeq \Sigma^{b_Y - p + 1} Y.$$

Proof. By Lemma 5.1, $\bar{A}_{p-1}^{\min} = S^{b_Y - p + 2}$. By Lemma 5.2 (4),

$$[Y \wedge \bar{A}^{\min}(Y)]_p \simeq \Sigma^{b_Y - p + 1} Y.$$

The result then follows from Lemma 5.2 (5). $\square$

Proof of Theorem 1.5. (1) and (2). Let

$$H_p^Y: \Omega Y \longrightarrow \Omega[\Sigma\Omega Y]_p$$

be the James-Hopf invariant. By Lemma 5.3,

$$\bar{A}_p^{\min}(Y) \simeq \Sigma^{b_Y - p + 1} Y.$$

Define the map H_p by the composite

$$\begin{array}{ccccc} \bar{A}^{\min}(Y) & \hookrightarrow & \Omega Y & \xrightarrow{H_p^Y} & \Omega([\Sigma\Omega Y]_p) \\ & \twoheadrightarrow & \Omega(\bar{A}_p^{\min}(Y)) & \twoheadrightarrow & \bar{A}^{\min}(\bar{A}_p^{\min}(Y)) = \bar{A}^{\min}(\Sigma^{b_Y - p + 1} Y). \end{array}$$

Let $\bar{E}(Y)$ be the homotopy fibre of the map H_p . Then there is a fibre sequence

$$\bar{E}(Y) \longrightarrow \bar{A}^{\min}(Y) \xrightarrow{H_p} \bar{A}^{\min}(\bar{A}_p^{\min}(Y)).$$

We compute the homology of $\bar{E}(Y)$. Let $V = \Sigma^{-1}\bar{H}_*(Y)$. According to Theorem 4.6,

$$E^0 H_{p*} = H_p: T(V) = E^0 H_*(\Omega Y) \longrightarrow T(V^{\otimes p}) = E^0 H_*(\Omega([\Sigma \Omega Y]_p))$$

is the usual algebraic James-Hopf map. Thus there is a commutative diagram

$$\begin{array}{ccc} T(V) & \xrightarrow{H_p} & T(V^{\otimes p}) \\ \downarrow & & \downarrow \\ E^0 H_*(\bar{A}^{\min}(Y)) = A^{\min}(V) & \xrightarrow{E^0 H_{p*}} & E^0 H_*(\bar{A}^{\min}(\bar{A}_p^{\min}(Y))) \end{array}$$

where H_p is the algebraic James-Hopf map. By Theorem 1.1 and Lemma 3.2, the primitives of $A^{\min}(V)$ is given by

$$PA^{\min}(V) = \bigoplus_{k=0}^{\infty} \bar{L}_p^k(V).$$

Let $\bar{L}$ be the sub Lie algebra of $L(V)$ generated by $L_p(V)$. Note that

$$\bar{L}_p^k(V) \subseteq \bar{L}$$

for $k \geq 1$. By [21, Theorem 1.1], the James-Hopf map

$$H_p|_{\bar{L}}: \bar{L} \longrightarrow PT(V^{\otimes p})$$

is a morphism of Lie algebras. Since

$$H_p|_{L_p}: \bar{L}_p \rightarrow V^{\otimes p}$$

is the canonical inclusion,

$$H_p(\bar{L}_p^{k+1}(V)) = \bar{L}_p^k(\bar{L}_p(V))$$

for each $k \geq 0$. Note that $\bar{L}_p(V) \subseteq A_p^{\min}(V)$ and, since $\dim V = p-1$,

$$\dim \bar{L}_p(V) = \dim A_p^{\min}(V) = p-1.$$

Thus $\bar{L}_p(V) = A_p^{\min}(V)$. It follows that

$$(5.3) \quad PE^0 H_{p*}: PE^0 H_*(\bar{A}^{\min}(Y)) \longrightarrow PE^0 H_*(\bar{A}^{\min}(\bar{A}_p^{\min}(Y)))$$

is onto with the kernel $L_p^0(V) = V$.

For any graded Hopf algebra A , let A^{ab} be the abelianization of A and let

$$\phi: A \longrightarrow A^{\text{ab}}$$

be the quotient map. Define $\bar{\phi}$ to be the composite

$$H_*(\bar{A}^{\min}(Y)) \hookrightarrow H_*(\Omega Y) \xrightarrow{\phi} H_*(\Omega Y)^{\text{ab}}.$$

Since $H_*(\Omega Y)$ is generated by odd dimensional elements, $H_*(\Omega Y)^{\text{ab}}$ is an exterior algebra. Moreover $E^0 H_*(\Omega Y)^{\text{ab}}$ is primitively generated exterior algebra. Thus

$$(5.4) \quad E^0 H_*(\Omega Y)^{\text{ab}} = E(V).$$

Consider the map

$$(5.5) \quad \theta: H_*(\bar{A}^{\min}(Y)) \begin{array}{c} \xrightarrow{\psi} \\ \xrightarrow{\bar{\phi} \otimes H_{p*}} \end{array} \begin{array}{c} H_*(\bar{A}^{\min}(Y)) \otimes H_*(\bar{A}^{\min}(Y)) \\ H_*(\Omega Y)^{\text{ab}} \otimes H_*(\bar{A}^{\min}(\bar{A}_p^{\min}(Y))) \end{array}.$$

From equations (5.3) and (5.4),

$$E^0 P\theta: PE^0 H_*(\bar{A}^{\min}(Y)) \longrightarrow PE^0(H_*(\Omega Y)^{\text{ab}} \otimes H_*(\bar{A}^{\min}(\bar{A}_p^{\min}(Y))))$$

is an isomorphism. Thus

$$E^0 \theta: E^0 H_*(\bar{A}^{\min}(Y)) \longrightarrow E^0(H_*(\Omega Y)^{\text{ab}} \otimes H_*(\bar{A}^{\min}(\bar{A}_p^{\min}(Y))))$$

is a monomorphism. Since both sides have the same Poincaré series, $E^0 \theta$ is an isomorphism and so θ is an isomorphism. Thus there is an algebra isomorphism on cohomology

$$H^*(\bar{A}^{\min}(Y)) \cong H^*(\bar{A}^{\min}(\bar{A}_p^{\min}(Y))) \otimes (H_*(\Omega Y)^{\text{ab}})^*.$$

By the Eilenberg-Moore spectral sequence, there is an epimorphism

$$H_*(\Omega Y)^{\text{ab}} \twoheadrightarrow H_*(\bar{E}(Y))$$

and so the Poincaré series

$$\chi(H_*(\bar{E}(Y))) \leq \frac{\chi(H_*(\bar{A}^{\min}(Y)))}{\chi(H_*(\bar{A}^{\min}(\bar{A}_p^{\min}(Y))))}.$$

By applying the Serre spectral sequence to the fibre sequence

$$\bar{E}(Y) \longrightarrow \bar{A}^{\min}(Y) \longrightarrow \bar{A}^{\min}(\bar{A}_p^{\min}(Y)),$$

we have the E^2 -terms given by

$$H_*(\bar{E}(Y)) \otimes H_*(\bar{A}^{\min}(\bar{A}_p^{\min}(Y)))$$

and so

$$\chi(H_*(\bar{A}^{\min}(Y))) \leq \chi(H_*(\bar{E}(Y))) \chi(H_*(\bar{A}^{\min}(\bar{A}_p^{\min}(Y)))).$$

It follows that

$$\chi(H_*(\bar{E}(Y))) = \frac{\chi(H_*(\bar{A}^{\min}(Y)))}{\chi(H_*(\bar{A}^{\min}(\bar{A}_p^{\min}(Y))))}$$

and so

$$H_*(\Omega Y)^{\text{ab}} \cong H_*(\bar{E}(Y)).$$

This finishes the proof of assertion (1) and (2).

(3). Let $\bar{Y}$ be the homotopy cofibre of the map $f: S^n \rightarrow Y$. Since f is a co- H -map, $\bar{Y}$ is a co- H -space. By the naturality of $\bar{A}^{\min}$, the pinch map $q: Y \rightarrow \bar{Y}$ induces a map

$$\bar{A}^{\min}(Y) \longrightarrow \bar{A}^{\min}(\bar{Y}).$$

Let $V' = \Sigma^{-1} \text{Im}(f_*: \bar{H}_*(S^n) \rightarrow \bar{H}_*(Y))$ and let $\bar{V} = \Sigma^{-1} \bar{H}_*(\bar{Y}) \cong V/V'$. Since $\dim \bar{V} = p - 2$,

$$(5.6) \quad E^0 H_*(\bar{A}^{\min}(\bar{Y})) = E(\bar{V})$$

by [16, Corollary 11.6]. From the proof of assertions (1) and (2), the composite

$$(5.7) \quad H_*(\bar{E}(Y)) \hookrightarrow H_*(\bar{A}^{\min}(Y)) \hookrightarrow H_*(\Omega Y) \twoheadrightarrow H_*(\Omega Y)^{\text{ab}}$$

is an isomorphism. Let g be the composite

$$\bar{E}(Y) \longrightarrow \bar{A}^{\min}(Y) \longrightarrow \bar{A}^{\min}(\bar{Y}).$$

Consider the commutative diagram

$$(5.8) \quad \begin{array}{ccccccc} H_*(\bar{E}(Y)) & \hookrightarrow & H_*(\bar{A}^{\min}(Y)) & \hookrightarrow & H_*(\Omega Y) & \twoheadrightarrow & H_*(\Omega Y)^{\text{ab}} \\ & \searrow \scriptstyle g_* & \downarrow & & \downarrow & & \downarrow \\ & & H_*(\bar{A}^{\min}(\bar{Y})) & \hookrightarrow & H_*(\Omega \bar{Y}) & \twoheadrightarrow & H_*(\Omega \bar{Y})^{\text{ab}}, \end{array}$$

where the composites in the top and bottom rows are isomorphisms. Thus

$$g_*: H_*(\bar{E}(Y)) \rightarrow H_*(\bar{A}^{\min}(Y))$$

is onto. By [12, Proposition 4.20], both $H_*(\Omega Y)^{\text{ab}}$ and $H_*(\Omega \bar{Y})^{\text{ab}}$ are primitively generated because they are commutative with trivial restricted maps. Thus

$$H_*(\Omega \bar{Y})^{\text{ab}} \cong E(\bar{V}) \text{ and } H_*(\Omega Y)^{\text{ab}} \cong E(V) \cong E(\bar{V}) \otimes E(V')$$

as Hopf algebras. From diagram 5.8 the map g_* induces a coalgebra decomposition

$$H_*(\bar{E}(Y)) \cong H_*(\bar{A}^{\min}(\bar{Y})) \otimes E(V').$$

It follows that the homotopy fibre of the map $g: \bar{E}(Y) \rightarrow \bar{A}^{\min}(\bar{Y})$ is S^{n-1} by using the Eilenberg-Moore spectral sequence. Thus there is a homotopy commutative diagram of fibre sequences

$$\begin{array}{ccccccc} \Omega \bar{A}^{\min}(\bar{A}_p^{\min}(Y)) & \xrightarrow{P_f} & S^{n-1} & \longrightarrow & B_f & \longrightarrow & \bar{A}^{\min}(\bar{A}_p^{\min}(Y)) \\ \parallel & & \downarrow & & \downarrow & & \parallel \\ \Omega \bar{A}^{\min}(\bar{A}_p^{\min}(Y)) & \xrightarrow{P} & \bar{E}(Y) & \xrightarrow{E} & \bar{A}^{\min}(Y) & \xrightarrow{H_p} & \bar{A}^{\min}(\bar{A}_p^{\min}(Y)) \\ & & \downarrow & & \downarrow & & \\ & & \bar{A}^{\min}(\bar{Y}) & = & \bar{A}^{\min}(\bar{Y}) & & \end{array}$$

and hence the result. $\square$

5.2. Self-maps of $\bar{A}^{\min}(Y)$. Let Y be a simply connected co- H -space of $(p-1)$ -cell complex with the cells in even dimensions. In this case, the integral homology of Y is torsion free and so we can take homology $H_*(Y)$ over p -local integers. Let f be any self-map of $\bar{A}^{\min}(Y)$. Let $V = \Sigma^{-1} \bar{H}_*(Y)$ with a basis $\{x_{l_1}, \dots, x_{l_{p-1}}\}$, where the degree $|x_{l_i}| = l_i$ with $l_1 \leq \dots \leq l_{p-1}$. By Theorem 1.5, there is a coalgebra isomorphism

$$H_*(\bar{A}^{\min}(Y)) \cong \bigotimes_{k=0}^{\infty} H_*(\bar{E}(\bar{A}_{p^k}^{\min}(Y))) \cong \bigotimes_{k=0}^{\infty} E(A_{p^k}^{\min}(V)) \cong A^{\min}(V).$$

Thus the primitives

$$PH_*(\bar{A}^{\min}(Y)) = \bigoplus_{k=0}^{\infty} A_{p^k}^{\min}(V)$$

and so on cohomology

$$QH^*(\bar{A}^{\min}(Y)) = \bigoplus_{k=0}^{\infty} A_{p^k}^{\min}(V)^*.$$

Note that $A_{p^k}^{\min}(V)$ has a basis in the form

$$w_{p^k,i} = [(x_{l_1} \wedge \cdots \wedge x_{l_{p-1}})^{\frac{p^k-1}{p-1}} x_{l_i}]$$

for $1 \leq i \leq p-1$. Let $f_{k,l}^*$ be the composite

$$A_{p^k}^{\min}(V)^* \hookrightarrow QH^*(\bar{A}^{\min}(Y)) \xrightarrow{f^*} QH^*(\bar{A}^{\min}(Y)) \xrightarrow{\text{proj.}} A_{p^k}^{\min}(V)^*.$$

Note that $|w_{p^k,p-1}| < |w_{p^{k'},1}|$ for $k < k'$. Thus

$$(5.9) \quad f_{k,l}^* = 0 \text{ for } k \neq l$$

and so

$$(5.10) \quad f^*(A_{p^k}^{\min}(V)^*) \subseteq A_{p^k}^{\min}(V)^*.$$

It follows that

$$f^*: A^{\min}(V)^* = H^*(\bar{A}^{\min}(Y)) \longrightarrow H^*(\bar{A}^{\min}(Y)) = A^{\min}(V)^*$$

is a graded map with respect to the tensor length. Since $\dim A_{p^{k-1}}^{\min}(V) = 1$, let $\deg^k(f)$ be the degree of the map

$$f^*: A_{p^{k-1}}^{\min}(V)^* \longrightarrow A_{p^{k-1}}^{\min}(V)^*.$$

Proposition 5.4. *Let $f: \bar{A}^{\min}(Y) \rightarrow \bar{A}^{\min}(Y)$ be any map. Then $\deg^k(f)$ over p -local integers is invertible if and only if*

$$f_*: H_j(\bar{A}^{\min}(Y); \mathbb{Z}/p) \longrightarrow H_j(\bar{A}^{\min}(Y); \mathbb{Z}/p)$$

is an isomorphism for $j < (l_1 + \cdots + l_{p-1}) \cdot \frac{p^k-1}{p-1} + l_1$.

Proof. The assertion follows from that

$$\deg^k(f) = \prod_{j=0}^{k-1} \det \left(f^*: A_{p^j}^{\min}(V)^* \longrightarrow A_{p^j}^{\min}(V)^* \right)$$

for cohomology with coefficients over p -local integers. $\square$

Since $\bar{A}^{\min}(Y)$ is an H -space, there is a product $f * g$ for any self maps f and g of $\bar{A}^{\min}(Y)$.

Proposition 5.5. *Let f and g be any self maps of $\bar{A}^{\min}(Y)$. Then*

$$\deg^k(f * g) = \prod_{j=0}^{k-1} \det \left(f_* + g_*: A_{p^j}^{\min}(V) \longrightarrow A_{p^j}^{\min}(V) \right).$$

Proof. Note that

$$(f * g)_*(w) = f_*(w) + g_*(w)$$

for any primitive element w . The assertion follows. $\square$

6. PROOF OF THEOREM 1.6

Lemma 6.1. *Let Y be a p -local simply connected co- H -space such that $H_{\text{odd}}(Y) = 0$ and $\dim \bar{H}_*(Y) = p-1$.*

- (1). *If Y admits a nontrivial decomposition, then the EHP fibration splits off.*
- (2). *If Y is atomic, then $\bar{E}(Y)$ is also atomic.*

Proof. (1). Suppose that $Y \simeq Y_1 \vee Y_2$ be a nontrivial decomposition. Since $\dim \bar{H}_*(Y_i) < p-1$, $H_*(\bar{A}^{\min}(Y_i)) = E(\Sigma^{-1}\bar{H}_*(Y_i))$. Let p_i be the composite

$$\bar{A}^{\min}(Y) \hookrightarrow \Omega Y \xrightarrow{\text{proj.}} \Omega Y_i \longrightarrow \bar{A}^{\min}(Y_i).$$

Then the composite

$$\bar{E}(Y) \xrightarrow{E} \bar{A}^{\min}(Y) \xrightarrow{(p_1, p_2)} \bar{A}^{\min}(Y_1) \times \bar{A}^{\min}(Y_2)$$

is a homotopy equivalence as it induces an isomorphism on homology. Assertion (1) follows.

(2). Let $f: \bar{E}(Y) \rightarrow \bar{E}(Y)$ be any self map inducing an isomorphism on the bottom cell. Let $V = \Sigma^{-1}\bar{H}_*(Y)$. By Theorem 1.5, $H_*(\bar{E}(Y)) = E(V)$ as coalgebras. Then

$$Pf_*: V = PE(V) \longrightarrow V = PE(V)$$

is an isomorphism on the bottom cell. From the suspension splitting theorem (Theorem 4.4),

$$\Sigma \bar{E}(Y) \simeq \bigvee_{n=1}^{p-1} \bar{A}_n^{\min}(Y)$$

with $\Sigma^{-1}H_*(\bar{A}_n^{\min}(Y)) = A^{\min}(V)$. Now the composite

$$g: Y = \bar{A}_1^{\min}(Y) \hookrightarrow \Sigma \bar{E}(Y) \xrightarrow{\Sigma f} \Sigma \bar{E}(Y) \simeq \bigvee_{n=1}^{p-1} \bar{A}_n^{\min}(Y) \longrightarrow \bar{A}_1^{\min}(Y) = Y$$

is a homotopy equivalence because on homology $\Sigma^{-1}g_* = Pf_*: V \rightarrow V$ is an isomorphism on the bottom cell. It follows that $Pf_*: V \rightarrow V$ is an isomorphism and so $f_*: E(V) \rightarrow E(V)$ is an isomorphism. Thus f is a homotopy equivalence and hence the result. $\square$

Let $H_*(\Omega Y)$ be filtered by the products of the augmentation ideal. For spaces X and Z such that $H_*(X)$ and $H_*(Z)$ are filtered, a map $f: X \rightarrow Z$ is called *filtered* if $f_*: H_*(X) \rightarrow H_*(Z)$ is a filtered map. A retract X of ΩY is called *filtered* if there is a filtered self map f of ΩY such that $X \simeq \text{hocolim}_f \Omega Y$. Note that for each filtered retract X of ΩY there is a filtration on $H_*(X)$ induced from the filtration of $H_*(\Omega Y)$. Consider all possible filtered retracts of ΩY . Divide them in two types: a filtered retract X of ΩY is called of *type A* if X contains the bottom cells of ΩY , otherwise it is called of *type B*.

Lemma 6.2. *For any p -local simply connected co- H -space Y , there exists a minimal filtered retract $X^{\min}(Y)$ of ΩY of type A such that: (1) $X^{\min}(Y)$ is a filtered retract of ΩY of type A and (2) any filtered retract of ΩY of type A is a filtered retract of $X(Y)$. Moreover there is a filtered decomposition*

$$\Omega Y \simeq X^{\min}(Y) \times \Omega Q.$$

Proof. Let $\mathcal{S}$ be the set of all filtered retract of ΩY of type A . Define a partial order on $\mathcal{S}$ by setting $X \leq X'$ if X is a filtered retract of X' . From $X_1, X_2 \in \mathcal{S}$, let f be the composite

$$\Omega Y \twoheadrightarrow X_1 \hookrightarrow \Omega Y \twoheadrightarrow X_2 \hookrightarrow \Omega Y.$$

Let $X_3 = \text{hocolim}_f \Omega Y$. Then X_3 is common filtered retract of X_1 and X_2 . Namely $X_3 \leq X_1$ and $X_3 \leq X_2$. By Zorn Lemma, $\mathcal{S}$ has the minimal element $X^{\min}(Y)$. Consider the commutative diagram

$$\begin{array}{ccccccc} \Omega \Sigma X^{\min}(Y) & \rightarrow & X^{\min}(Y) & \rightarrow & \Sigma X^{\min}(Y) \wedge X^{\min}(Y) & \xrightarrow{H} & \Sigma X^{\min}(Y) \\ \uparrow \Omega j & & \parallel & & \uparrow & \text{pull} & \uparrow j \\ \Omega Y & \xrightarrow{\partial} & X^{\min}(Y) & \longrightarrow & Q & \longrightarrow & Y \end{array}$$

where j is the adjoint map of the retraction $\Omega Y \rightarrow X^{\min}(Y)$. Then the composite

$$X^{\min}(Y) \longrightarrow \Omega Y \xrightarrow{\partial} X^{\min}(Y)$$

is a homotopy equivalence by the minimal assumption of $X^{\min}(Y)$. It follows that there is a filtered decomposition

$$\Omega Y \simeq X^{\min}(Y) \times \Omega Q$$

and hence the result. $\square$

Proof of Theorem 1.6. (1) $\implies$ (2). If EHP fibration splits off, then $\bar{E}(Y)$ is a retract of $\bar{A}^{\min}(Y)$. Note that $\bar{A}^{\min}(Y)$ is an H -space as it is a retract of ΩY . Thus $\bar{E}(Y)$ is an H -space.

(2) $\implies$ (3). From the suspension splitting theorem (Theorem 4.4),

$$\Sigma \bar{A}^{\min}(Y) \simeq \bigvee_{n=1}^{\infty} \bar{A}_n^{\min}(Y)$$

with $\bar{A}_1^{\min}(Y) = Y$ and $\Sigma^{-1} \bar{H}_*(\bar{A}_n^{\min}(Y)) = A^{\min}(\Sigma^{-1} \bar{H}_*(Y))$. The composite

$$\Sigma \bar{E}(Y) \longrightarrow \Sigma \bar{A}^{\min}(Y) \xrightarrow{\text{proj.}} \bigvee_{n=1}^{p-1} \bar{A}_n^{\min}(Y)$$

is a homotopy equivalence because $H_*(\bar{E}(Y)) \rightarrow H_*(\bar{A}^{\min}(Y))$ is a monomorphism and the connectivity of $\bigvee_{n=p}^{\infty} \bar{A}_n^{\min}(Y)$ is greater than the dimension of $\Sigma \bar{E}(Y)$. Thus Y is a retract of $\Sigma \bar{E}(Y)$. By Theorem 1.5, $H^*(\bar{E}(Y)) = E(\Sigma^{-1} \bar{H}^*(Y))$ and so $\bar{E}(Y)$ is an H -space with Y as a retractile generating complex.

(3) $\implies$ (1). Let X be any H -space having Y as a retractile generating complex. Since Y is a retract of X . Let $s: Y \rightarrow \Sigma X$ be an inclusion and let $r: \Sigma X \rightarrow Y$ be the retraction such that $H^*(X)$ is generated by $M = r^*(\Sigma^{-1} \bar{H}_*(Y))$ with $M \cong QH^*(X)$. Let $V = \Sigma^{-1} \bar{H}_*(Y)$. Since $H_{\text{odd}}(Y) = 0$, $V_{\text{even}} = 0$. Note that $H_*(X)$ is a quasi-Hopf algebra as X is an H -space. Thus $H^*(X)$ is the exterior algebra generated by V^* by the Borel Theorem [12, Theorem 7.11]. There is a homotopy

commutative diagram of fibre sequences

$$(6.1) \quad \begin{array}{ccccccc} \Omega\Sigma X & \xrightarrow{\tilde{\partial}} & X & \longrightarrow & \Sigma X \wedge X & \xrightarrow{H} & \Sigma X \\ \uparrow \Omega s & & \parallel & & \uparrow \text{pull} & & \uparrow s \\ \Omega Y & \xrightarrow{\partial} & X & \longrightarrow & B & \longrightarrow & Y, \end{array}$$

where $H: \Sigma X \wedge X \rightarrow \Sigma X$ is the Hopf fibration for the H -space X . Let $r': X \rightarrow \Omega Y$ be the adjoint map of r . Since the dimension of X less than the connectivity of $\bar{A}^{\min}(\Sigma^{b_Y-p+1}(Y))$, the composite

$$X \xrightarrow{r'} \Omega Y \longrightarrow \bar{A}^{\min}(Y)$$

lifts to the fibre $\bar{E}(Y)$ of the EHP-fibration. Let $r'': X \rightarrow \bar{E}(Y)$ be a lifting of the above composite. Consider the composite

$$(6.2) \quad \begin{array}{ccccccccc} \phi: \Omega Y & \xrightarrow{\partial} & X & \xrightarrow{r''} & \bar{E}(Y) & \xrightarrow{E} & \bar{A}^{\min}(Y) & \hookrightarrow & \Omega Y \\ \uparrow j & & \uparrow j & & \uparrow j & & \uparrow j & & \uparrow j \\ \bigvee_{\alpha} S^n & \xlongequal{\quad} & \bigvee_{\alpha} S^n & \xlongequal{\quad} & \bigvee_{\alpha} S^n & \xlongequal{\quad} & \bigvee_{\alpha} S^n & \xlongequal{\quad} & \bigvee_{\alpha} S^n, \end{array}$$

where j is the inclusion of the bottom cells. Let

$$\bar{X} = \text{hocolim}_{\phi} \Omega Y.$$

Then $\bar{X}$ is a common retract of X , $\bar{E}(Y)$, $\bar{A}^{\min}(Y)$ and ΩY , which contains the bottom cell. Since Y is atomic, $\bar{E}(Y)$ is also atomic by Lemma 6.1. Thus the retraction

$$\bar{E}(Y) \xrightarrow{E} \bar{A}^{\min}(Y) \hookrightarrow \Omega Y \longrightarrow \text{hocolim}_{\phi} \Omega Y = \bar{X}$$

is a homotopy equivalence. It follows that the EHP-sequence

$$\bar{E}(Y) \xrightarrow{E} \bar{A}^{\min}(Y) \xrightarrow{H_p} \bar{A}^{\min}(\Sigma^{b_Y-p+1}Y)$$

admits a retraction $\bar{A}^{\min}(Y) \rightarrow \bar{E}(Y)$ and so the EHP fibration splits off.

Now (1) $\implies$ (4) and (4) $\implies$ (5) are obvious. (5) $\implies$ (6) follows from the homotopy commutative diagram

$$\begin{array}{ccccccc} \Omega \bar{A}^{\min}(\Sigma^{b_Y-p+1}Y) & \xrightarrow{P} & \bar{E}(Y) & \xrightarrow{E} & \bar{A}^{\min}(Y) & \xrightarrow{H_p} & \bar{A}^{\min}(\Sigma^{b_Y-p+1}Y) \\ \uparrow & & \uparrow & & \uparrow g & & \uparrow \\ \Sigma^{b_Y-p-1}Y & \longrightarrow & * & \longrightarrow & \Sigma^{b_Y-p}Y & \xlongequal{\quad} & \Sigma^{b_Y-p}Y, \end{array}$$

where the bottom row is the cofibre sequence.

(6) $\implies$ (3). Consider the commutative diagram

$$\begin{array}{ccccccc}
 \bar{A}^{\min}(\Sigma^{b_Y-p+1}Y) & \hookrightarrow & \Omega\Sigma^{b_Y-p+1}Y & \xrightarrow{\tilde{g}} & \Omega Y & \longrightarrow & \bar{A}^{\min}(Y) \\
 & & \uparrow & & \uparrow & \nearrow & \\
 & & \Sigma^{b_Y-p}Y & \xrightarrow{g} & \bar{A}^{\min}(Y), & &
 \end{array}$$

where $\tilde{g}$ is the H -map induced by g . Let $\bigvee_{\alpha} S^q$ be the bottom cells of $\Sigma^{b_Y-p}Y$. Observe that

$$(6.3) \quad H_{p*}: H_q(\bar{A}^{\min}(Y)) \xrightarrow{\cong} H_q(\bar{A}^{\min}(\Sigma^{b_Y-p+1}Y)).$$

Thus there is a commutative diagram

$$\begin{array}{ccccccccc}
 \theta: \Omega Y & \longrightarrow & \bar{A}^{\min}(Y) & \xrightarrow{H_p} & \bar{A}^{\min}(\Sigma^{b_Y-p+1}Y) & \hookrightarrow & \Omega\Sigma^{b_Y-p+1}Y & \xrightarrow{\tilde{g}} & \Omega Y \\
 \uparrow & & \uparrow & & \uparrow & & \uparrow & & \uparrow \\
 \bigvee_{\alpha} S^q & \xlongequal{\quad} & \bigvee_{\alpha} S^q & \xlongequal{\quad} & \bigvee_{\alpha} S^q & \xlongequal{\quad} & \bigvee_{\alpha} S^q & \xlongequal{\quad} & \bigvee_{\alpha} S^q, \\
 & & g|_{\bigvee_{\alpha} S^q} & & j & & j & &
 \end{array}
 \quad (6.4)$$

where j is the inclusion of the bottom cells.

Let $H_*(\Omega Y)$ be filtered by the products of the augmentation ideal. Then $H_*(\bar{A}(Y))$ has the induced filtration. Observe that

$$g_*(\bar{H}_*(\Sigma^{b_Y-p}Y)) \subseteq I^p H_*(\bar{A}^{\min}(Y)) \subseteq I^p H_*(\Omega Y)$$

because the top dimension of $H_*(\bar{E}(Y)) = \bigoplus_{n=1}^{p-1} \bar{A}^{\min}(\Sigma^{-1}Y)$ is less than the connectivity of $\Sigma^{b_Y-p}Y$. It follows that

$$\tilde{g}_*: H_*(\Omega\Sigma^{b_Y-p+1}Y) = H_*(J(\Sigma^{b_Y-p}Y)) \longrightarrow H_*(\Omega Y)$$

preserves the filtration. Thus θ is a filtered map because the other factors of θ are filtration-preserving maps on homology. Let

$$Z = \text{hocolim}_{\theta} \Omega Y$$

be the homotopy colimit. Then Z is a common filtered retract of ΩY , $\bar{A}^{\min}(Y)$, $\bar{A}^{\min}(\Sigma^{b_Y-p+1}Y)$, and $\Omega\Sigma^{b_Y-p+1}Y$. By Lemma 4.3, ΣZ is a common graded retract of $\Sigma\Omega Y$, $\Sigma\bar{A}^{\min}(Y)$, $\Sigma\bar{A}^{\min}(\Sigma^{b_Y-p+1}Y)$, and $\Sigma\Omega\Sigma^{b_Y-p+1}Y$. Thus $[\Sigma Z]_p$ is a retract of

$$[\Sigma\bar{A}^{\min}(\Sigma^{b_Y-p+1}Y)]_p = \Sigma^{b_Y-p+1}Y$$

containing the bottom cells by diagram (6.4). It follows that

$$(6.5) \quad [\Sigma Z]_p = [\Sigma\bar{A}^{\min}(\Sigma^{b_Y-p+1}Y)]_p = \Sigma^{b_Y-p+1}Y$$

because $\Sigma^{b_Y-p+1}Y$ is atomic. Thus Z contains $\Sigma^{b_Y-p}Y$ as the bottom piece.

Consider the filtered decomposition

$$\Omega Y \simeq \bar{B}^{\max}(Y) \times \bar{A}^{\min}(Y).$$

Since Z is a filtered retract of $\bar{A}^{\min}(Y)$, there is a further decomposition

$$\Omega Y \simeq \bar{B}^{\max}(Y) \times Z \times A',$$

where A' is a filtered retract of ΩY of type A . By Lemma 6.2, A' admits a further filtered decomposition

$$A' \simeq B' \times X^{\min}(Y).$$

This gives a filtered decomposition

$$(6.6) \quad \Omega Y \simeq \bar{B}^{\max}(Y) \times Z \times B' \times X^{\min}(Y)$$

and so the space Q given in Lemma 6.2 has the property that

$$(6.7) \quad \Omega Q \simeq \bar{B}^{\max}(Y) \times Z \times B'.$$

On homology

$$(6.8) \quad PE^0 H_*(Z) \oplus PE^0 H_*(\bar{B}^{\max}(Y)) \subseteq PE^0 H_*(\Omega Q).$$

Let $V = \Sigma^{-1} \bar{H}_*(Y)$. From equation (6.5),

$$(6.9) \quad \bar{L}_p(V) = A_p^{\min}(V) = \Sigma^{-1} \bar{H}_*([\Sigma \bar{A}^{\min}(\Sigma^{b_Y - p + 1} Y)]_p) \subseteq PE^0(\Omega Q).$$

By equation (6.8),

$$\bigoplus_{n=2}^{p-1} Q_n^{\max}(V) \oplus \bar{L}_p(V) \subseteq PH_*(\Omega Q).$$

Since $H_*(\Omega Q)$ is a Hopf algebra, the sub Hopf algebra

$$(6.10) \quad B = \langle \bigoplus_{n=2}^{p-1} Q_n^{\max}(V) \oplus \bar{L}_p(V) \rangle \subseteq H_*(\Omega Q).$$

By Lemma 3.2, $B = B^{[1]}(V)$ with a short exact sequence of Hopf algebras

$$B \hookrightarrow T(V) \twoheadrightarrow E(V).$$

It follows that the Poincaré series

$$\chi(H_*(\Omega Q)) \geq \chi(B) = \frac{\chi(T(V))}{\chi(E(V))}.$$

From the decomposition

$$\Omega Y \simeq \Omega Q \times X^{\min}(Y),$$

the Poincaré series

$$\chi(H_*(X^{\min}(Y))) \leq \chi(E(V)).$$

Since $X^{\min}(Y)$ is filtered retract of ΩY , $[\Sigma X^{\min}(Y)]_1$ is a retract of $[\Sigma \Omega Y]_1 = Y$. Because Y is atomic, $[\Sigma X^{\min}(Y)]_1 \simeq Y$. It follows that

$$V \subseteq E^0 H_*(X^{\min}(Y)).$$

Since $X^{\min}(Y)$ is an H -space, $E^0 H_*(X^{\min}(Y))$ is a Hopf algebra and so

$$\chi(H_*(X^{\min}(Y))) = \chi(E^0 H_*(X^{\min}(Y))) \geq \chi(E(V)).$$

Thus $\chi(H_*(X^{\min}(Y))) = \chi(E(V))$ and so $H_*(X^{\min}(Y)) = E(V)$ as coalgebras. It follows that $X^{\min}(Y)$ is an H -space having Y as a retractile generating complex and hence statement (3). The proof is finished. $\square$

REFERENCES

- [1] F. Cohen, *On combinatorial group theory in homotopy*, Contemp. Math. **188** (1995), 57-63.
- [2] F. R. Cohen, J. C. Moore and J. A. Neisendorfer, *Torsion in homotopy groups*, Ann. of Math. (2) **109** (1979), 121-168.
- [3] G. Cooke, J. R. Harper and A. Zabrodsky, *Torsion free mod p H -spaces of low rank*, Topology **18** (1979), 349-359.
- [4] F. R. Cohen, J. Neisendorfer, *A construction of p -local H -spaces*, Algebraic Topology Lec. Notes in Math. **1051** (1982), Springer, 351-359.
- [5] Curtis and Reiner, *Methods of representation theory, Vol. I*, Pure and Applied Mathematics Wiley (1981).
- [6] J. Grbic, S. Theriault and J. Wu, *The suspension splitting of retracts of the loops on co- H -spaces and Hopf invariants*, preprint.
- [7] Jelena Grbic and Jie Wu, *Natural transformations of tensor algebras and representations of combinatorial groups*, Algebraic and Geometric Topology **6** (2006), 2189-2228.
- [8] Jelena Grbic and Jie Wu, *Applications of combinatorial groups to the Hopf invariants and the exponent problem*, Algebraic and Geometric Topology **6** (2006), 2229-2255.
- [9] Jelena Grbic, Paul Selick and Jie Wu, *The decomposition of the loop space of the mod 2 Moore space*, Algebraic and Geometric Topology **8** (2008) 945-951.
- [10] I. M. James, *Reduced product spaces*, Ann. Math. **62** (1953), 170-197.
- [11] J. Y. Li and J. Wu, *On natural coalgebra-split sub Hopf algebras of tensor algebras*, preprint.
- [12] J. Milnor and J. Moore, *On the structure of Hopf Algebras*, Ann. Math. **81** (1965), 211-264.
- [13] I. Schur, *Über eine Klasse von Matrizen, die sich einer gegebenen Matrix zuordnen lassen (1901)*, in I. Schur, Gesammelte Abhandlungen I, 1-70, Springer, Berlin, 1973.
- [14] Paul Selick, Stephen Theriault and Jie Wu, *Functorial decompositions of looped coassociative co- H spaces*, Canadian J. Math. **58** (2006), 877-896.
- [15] Paul Selick, Stephen Theriault and Jie Wu, *decompositions of looped co- H -spaces*, preprint.
- [16] Paul Selick and Jie Wu, *On natural decompositions of loop suspensions and natural coalgebra decompositions of tensor algebras*, Memoirs AMS **148** (2000), No. 701.
- [17] Paul Selick and Jie Wu, *The functor $A^{\min}$ on p -local spaces*, Math. Z. **253** (2006), 435-451.
- [18] P. S. Selick and J. Wu, *Some calculations of $\mathrm{Lie}^{\max}(n)$ for low n* , J. Pure. Appl. Algebra to appear.
- [19] Stephen D. Theriault, *Homotopy decompositions involving the loops of coassociative Co- H spaces*, Canad. J. Math. **55** (2003), 181-203.
- [20] H. Toda, *Composition methods in homotopy groups of spheres*, Annals of Mathematics Studies **49** (1962), Princeton University Press, Princeton, N.J., v+193 pp.
- [21] J. Wu, *On combinatorial calculations of the James-Hopf maps*, Topology **37** (1998), 1011-1023.
- [22] J. Wu, *Homotopy theory of the suspensions of the projective plane*, Memoirs AMS **162** (2003), No. 769.
- [23] J. Wu, *On maps from loop suspensions to loop spaces and the shuffle relations on the Cohen groups*, Memoirs AMS, **180** (2006), No. 851.

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119260,
REPUBLIC OF SINGAPORE

E-mail address: matwujie@math.nus.edu.sg

URL: <http://www.math.nus.edu.sg/~matwujie>